

Enhancer release and retargeting activates disease-susceptibility genes

<https://doi.org/10.1038/s41586-021-03577-1>

Received: 14 June 2018

Accepted: 23 April 2021

Published online: 26 May 2021



Soohwan Oh¹, Jiaofang Shao^{2,8}, Joydeep Mitra^{3,8}, Feng Xiong², Matteo D'Antonio⁴, Ruoyu Wang^{2,5}, Ivan Garcia-Bassets⁶, Qi Ma¹, Xiaoyu Zhu², Joo-Hyung Lee², Sreejith J. Nair¹, Feng Yang¹, Kenneth Ohgi¹, Kelly A. Frazer^{4,7}, Zhengdong D. Zhang⁵, Wenbo Li^{2,5}✉ & Michael G. Rosenfeld¹✉

The functional engagement between an enhancer and its target promoter ensures precise gene transcription¹. Understanding the basis of promoter choice by enhancers has important implications for health and disease. Here we report that functional loss of a preferred promoter can release its partner enhancer to loop to and activate an alternative promoter (or alternative promoters) in the neighbourhood. We refer to this target-switching process as ‘enhancer release and retargeting’. Genetic deletion, motif perturbation or mutation, and dCas9-mediated CTCF tethering reveal that promoter choice by an enhancer can be determined by the binding of CTCF at promoters, in a cohesin-dependent manner—consistent with a model of ‘enhancer scanning’ inside the contact domain. Promoter-associated CTCF shows a lower affinity than that at chromatin domain boundaries and often lacks a preferred motif orientation or a partnering CTCF at the cognate enhancer, suggesting properties distinct from boundary CTCF. Analyses of cancer mutations, data from the GTEx project and risk loci from genome-wide association studies, together with a focused CRISPR interference screen, reveal that enhancer release and retargeting represents an overlooked mechanism that underlies the activation of disease-susceptibility genes, as exemplified by a risk locus for Parkinson's disease (*NUCKS1–RAB7L1*) and three loci associated with cancer (*CLPTMIL–TERT*, *ZCCHC7–PAX5* and *PVT1–MYC*).

High-resolution data of chromatin architecture¹ indicate that enhancers and promoters form complex patterns of finer-scale loops (around 10–50 kb)²; however, functionally, most enhancers regulate one gene within a short distance³. What dictates the specificity and plasticity of enhancer(s) and promoter(s) engagement to license productive gene transcription remains unclear⁴.

Promoter loss activates partner enhancers

In MCF-7 cells—a well-established model of enhancer-dependent promoter activation⁵—treatment with 17- β -oestradiol causes the rapid activation of enhancer RNAs (eRNAs), which precedes the activation of target gene promoters⁶ (Extended Data Fig. 1a). To understand enhancer–promoter (E–P) choice, we selected four ligand-induced E–P pairs that exhibit looping by 4C-seq (Extended Data Fig. 1b) and chromatin interaction analysis with paired-end tag sequencing (ChIA-PET)⁷, and deleted each of these promoters encompassing transcription start sites (TSSs) (Supplementary Table 1, Supplementary Figs. 2, 4). In cells in which the *TFF1* promoter (*TFF1p*) was deleted (*TFF1p-KO* cells), the expression of *TFF1* mRNA was extremely reduced, but that of *TFF1* eRNA increased markedly (around three- to fivefold); genes

on other chromosomes (for example, *FOXCI*) exhibited no significant alterations (Fig. 1a). Results from more than 10 independent clones and using a second pair of guide RNAs (gRNAs) (Extended Data Fig. 1c, d) confirmed these findings. Similarly, deletion of the *P2RY2*, *KCNKS* and *PCR* promoters led to a reduction in the expression of their cognate genes but a significant increase in cognate eRNA transcription (Extended Data Fig. 2a, d, g). The *TFF1* enhancer (*TFF1e*)—but not control enhancers—showed an increase in the binding of RNA polymerase II (Pol II) and p300 in *TFF1p-KO* cells (Extended Data Fig. 1e). These results support the notion that ligand-induced enhancers are activated temporally before promoters and initiate promoter choice.

Results from cohesin knockdown experiments support this premise. Knockdown of *RAD21* (which encodes a component of the cohesin complex) disrupted E–P loops in the loci we tested (Extended Data Figs. 1b, 3a), and concomitantly, GRO-seq and quantitative PCR with reverse transcription (qRT–PCR) showed that genes were no longer effectively induced by treatment with 17- β -oestradiol, whereas eRNAs were significantly upregulated (Fig. 1b, Extended Data Fig. 3b–f). No significant alteration of methylation of histone H3 at K4 (H3K4) was detected (Extended Data Fig. 3g).

¹Howard Hughes Medical Institute, Department and School of Medicine, University of California San Diego, La Jolla, CA, USA. ²Department of Biochemistry and Molecular Biology, McGovern Medical School, University of Texas Health Science Center, Houston, TX, USA. ³Department of Genetics, Albert Einstein College of Medicine, New York, NY, USA. ⁴Department of Pediatrics, School of Medicine, University of California San Diego, La Jolla, CA, USA. ⁵Graduate School of Biomedical Sciences, University of Texas MD Anderson Cancer Center and UTHealth, Houston, TX, USA. ⁶Department of Medicine, School of Medicine, University of California San Diego, La Jolla, CA, USA. ⁷Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA. ⁸These authors contributed equally: Jiaofang Shao, Joydeep Mitra. ✉e-mail: Wenbo.Li@uth.tmc.edu; mrosenfeld@health.ucsd.edu




Fig. 1 | Promoter deletion causes ERR. **a**, Box plots from qRT-PCR in wild-type (WT) ($n = 6$) and *TFF1p-KO* ($n = 14$) independent cell clones, showing the expression of *TFF1* mRNA (*TFF1m*), *TFF1* enhancer RNA (*TFF1e*) and *FOXC1m* relative to *GAPDH*. The left diagram shows the *TFF1p* deletion. **b**, Metagene plots showing GRO-seq signals from downregulated promoters or super-activated eRNAs after small interfering RNA (siRNA)-mediated knockdown of *RAD21* (si*RAD21*). E_2 , 17- β -oestradiol. **c**, Hi-C map from GM12878 cells⁸ (top, by Juicebox); CTCF ChIA-PET in MCF-7 (from Gene Expression Omnibus (GEO) accession GSM970215) (middle); and indicated ChIP-seq tracks showing the chromosomal topology around the *TFF1* locus (bottom). Yellow triangles in the Hi-C map denote contact domains, which are numbered for simplicity. The

domain containing *TFF1* is highlighted in yellow. **d**, Box plots from qRT-PCR in wild-type ($n = 6$) versus *TFF1p-KO* ($n = 14$) cell clones, showing the expression of all mRNAs (relative to *GAPDH*) in the domain hosting the *TFF1*-E-P pair. **e**, 4C-seq in wild-type versus *TFF1p-KO* cells, based on 'viewpoints' (asterisks) centred on *TFF3p* (yellow) or *TFF1e* (pink). **f**, Browser tracks showing GRO-seq and H3K4me3 ChIP-seq in the *TFF1*-*TFF3* region in wild-type versus *TFF1p-KO* cells; arrowheads point to gained H3K4me3 on *TFF3p*. The box plot centre lines represent medians; box limits indicate the 25th and 75th percentiles; and whiskers extend 1.5 times the interquartile range (IQR) from the 25th and 75th percentiles. ** $P < 0.01$, *** $P < 0.001$, NS, not significant, two-tailed Student's *t*-test.

Promoters determine E-P specificity

TFF1 and four other genes are located in an approximately 150-kb contact domain⁸ (Fig. 1c), adjacent to several other discernible domains. After deletion of *TFF1p*, the expression of *TFF3* mRNA increased by more than 18-fold and the other three genes were also upregulated, albeit to a lesser extent (Fig. 1d). By contrast, genes in adjacent domains showed minimal changes in expression (Extended Data Fig. 4a–c). 4C-seq revealed an increased interaction between *TFF1e* and the *TFF3* promoter (*TFF3p*) after knockout of *TFF1p* (Fig. 1e). Levels of the promoter histone mark H3K4me3 increased on *TFF3p* (Fig. 1f), accompanying an increase in the binding of Pol II and p300 (Extended Data Fig. 4d). The new *TFF1e*–*TFF3p* contact still required cohesin, because knockdown of *RAD21* still led to an increase in the level of *TFF1e* eRNA, but a reduction in the level of *TFF3* mRNA (Extended Data Fig. 4e).

Deletion of *TFF1e* in *TFF1p-KO* cells, generating a double knockout (*TFF1e/p-DKO*; Supplementary Table 1, Supplementary Fig. 3), abolished the abnormal activation of all promoter targets in the domain (Extended Data Fig. 4f). This indicated that *TFF1e* drove the activation of alternative promoters after knockout of *TFF1p*. By contrast, deletion of *TFF1e* alone, although reducing the levels of *TFF1* mRNA, did not activate other genes (Extended Data Fig. 4f), indicating that the observed phenomenon—that is, super-activation of engaged enhancers and activation of alternative neighbouring promoters—was due to specific chromatin changes after promoter loss. Deletion of three other promoters tested had similar effects (Extended Data Fig. 2a, b, d, g, h). Together, these results suggest that the functional loss of cognate promoters 'releases' their enhancers to interact with and activate one

or more gene promoters in the neighbourhood. We refer to this process as 'enhancer release and retargeting' (ERR).

Promoter CTCF determines E-P engagement

Our identification of the process of ERR provided an opportunity to understand the specificity and plasticity of functional E-P engagement and promoter choice⁴. We found that all promoters that we deleted contain a CTCF-binding site (Extended Data Fig. 2c, f, i). Promoters that were functionally engaged in E-P looping, as suggested by their downregulation after knockdown of *RAD21*, showed a higher level of CTCF binding than random promoters (Extended Data Fig. 5a–c). Because CTCF is crucial for the formation of topologically associated domains (TADs)⁹, we hypothesized that promoter CTCF might have a role in determining finer-scale E-P choice and engagement.

Overall, promoters exhibit a higher CTCF binding affinity than enhancers, although the CTCF binding affinity at promoters is considerably weaker than that at domain boundaries (Fig. 2a). Around 30% of promoters in MCF-7 cells and around 20% in GM12878 cells exhibited CTCF binding (Extended Data Fig. 6a, b). CTCF binding was more common for promoters near a putative enhancer or super-enhancer¹⁰ (Extended Data Fig. 6c, d). Detectable CTCF binding occurs in around 60% of promoters of highly transcribed genes in MCF-7 cells (Fig. 2b). We experimentally examined the role of promoter-bound CTCF in *TFF1p*, which contains a CTCF motif and chromatin immunoprecipitation followed by sequencing (ChIP-seq) peak (Supplementary Table 1, Supplementary Fig. 2). Deletion of this CTCF peak (*TFF1p-ΔCTCF*) fully reproduced the effects of deletion of the entire *TFF1p* (Fig. 2c, e). Similar




Fig. 2 | CTCF binding at promoters is important for their functional engagement with enhancers. **a**, A normalized coverage plot showing differential CTCF binding at enhancers, promoters, and domain boundaries in MCF-7 cells. **b**, Bar graphs showing the numbers of promoters bearing CTCF ChIP-seq peaks for 1,000 promoters with high, medium or low levels of transcription (by GRO-seq) in MCF-7 cells. **** $P < 0.0001$, two-sided Fisher's exact test. **c**, qRT-PCR showing expression levels of the indicated targets in wild-type cells, *TFF1p*-KO cells or cells with deletion of the CTCF peak (Δ CTCF) or the ER α -binding site (Δ ERE). Each bar represents an independent cell clone ($n = 3$ biological replicates). **d**, qRT-PCR results showing altered *TFF1e*, *TFF1m* and *TFF3m* levels in cells in which the *TFF1p* CTCF motif was disrupted ($n = 3$

biological replicates). **e**, Browser screenshots showing hierarchical CTCF and cohesin (RAD21) binding at several gene promoters. CTCF peak (Δ CTCF) or ER α -binding site (Δ ERE) deletions are indicated. **f**, Diagram (top) and qRT-PCR results (bottom) showing the expression levels of *TFF1e* and several mRNAs in WT, *TFF1p*-KO and *TFF1p*/*TFF3p*-DKO cells ($n = 3$ biological replicates). **g**, Diagram (left) and qRT-PCR results (right) showing the relative expression levels of *TFF3m* and *TFF1m*, illustrating the strategy and effects of CTCF tethering ($n = 3$ biological replicates). Cells expressing dCas9-10xGCN4 and scFV-CTCF (wild type or Y226A/F228A mutant) were used. Data in **c**, **d**, **f**, **g** show mean $\pm$ s.d.; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's *t*-test.

changes were observed by disrupting the CTCF motif (Fig. 2d), but not by deleting oestrogen receptor recognition elements (EREs) (Fig. 2c, e).

TFF3p also contains a CTCF site, with an affinity lower than that of *TFF1p* but higher than that of *TFF2p* (Fig. 2e) or *UBASH3Ap*. Notably, deletion of *TFF3p* in *TFF1p*-KO cells (*TFF1p*/*TFF3p*-DKO; Supplementary Fig. 3) further increased the mRNA expression of *TFF2* and *UBASH3A* (Fig. 2f), suggesting that *TFF1e* is once again 'released' from its alternative target *TFF3p* to retarget other promoters in the domain. Tethering of CTCF to *TFF3p* (mediated by dCas9 and the SunTag system; see Methods) increased the expression of *TFF3* and reduced that of *TFF1*, whereas tethering of a CTCF mutant (Y226A/F228A)¹¹ defective for cohesin interaction had no effect (Fig. 2g, Extended Data Fig. 6e, f). On the basis of these data, all genes in the hosting domain appear to have the potential to be activated by *TFF1e*, suggesting that an enhancer scans the entire domain¹², with the binding of CTCF on candidate promoters serving as a component that determines the 'preferred' engagement by the enhancer (Extended Data Fig. 7).

CTCF motifs are preferentially convergent at TAD and loop domain boundaries^{8,9}, and for specific E-P loops¹³. ChIA-PET data showed that some RAD21-dependent promoters ($n = 148$) and super-activated enhancers after knockdown of *RAD21* ($n = 157$) exhibit MCF-7-specific loops (Extended Data Fig. 8a), which we considered to be functionally engaged E-P pairs. Of these pairs, around two-thirds have a CTCF peak on promoters, but only around 10% exhibit CTCF binding on both enhancer and promoter; there is no obvious preference of motif orientation for CTCF or for FOXA1 (Extended Data Fig. 8b,c). Of the four

E-P pairs studied above, all promoters exhibit CTCF binding, but only *TFF1e* contains a canonical CTCF motif and a peak. The CTCF motifs for *TFF1e*-P pair are not convergent (Extended Data Fig. 8d, Supplementary Table 2). Deletion of the *TFF1e* CTCF peak caused no discernible change of *TFF1* eRNA or mRNA (Extended Data Fig. 8e). These data indicate that, for functional E-P engagement, CTCF binding at promoters is required for at least a subset of promoters, whereas its presence at enhancers and convergent motif orientation are largely dispensable.

ERR-like events activate cancer genes

Cancer-associated single nucleotide polymorphisms (SNPs) are found in the *TFF1* promoter¹⁴. We investigated whether ERR could function as an overlooked mechanism that underlies disease-associated genetic alterations of promoters (that is, variations, mutations and deletions) (Fig. 3a). Analysis of data from the International Cancer Genome Consortium (ICGC) showed that each cancer type contains a large number of mutations near gene TSSs (Extended Data Fig. 9a), consistent with previous studies^{15,16}. We looked for genetic changes in other promoters that are located at a ± 200 kb distance from annotated oncogene promoters (OPs, $n = 315$), which we dubbed oncogene-neighbouring promoters (ONPs, $n = 1,693$) (Methods, Fig. 3a, Supplementary Figs. 6, 7). Overall, we found a moderate but significant enrichment of mutations in, or deletions of, OPs and ONPs, compared with the genome-wide average of random promoters (Extended Data Fig. 9b). Focusing on ONPs (Fig. 3a), we identified mutational hotspots at the promoter




Fig. 3 | ERR is a potentially common paradigm underlying oncogene activation. **a**, A model of ERR, proposing that noncoding mutations or variations of non-disease-causing promoters can deregulate the expression of disease-associated genes in the neighbourhood. **b**, qRT-PCR results showing altered levels of *TFF1e*, *TFF1m* and *TFF3m* after CRISPRi-mediated inhibition of *TFF1p* (iTFF1p) compared to a control (iCTL) ($n = 3$ biological replicates). The top diagram shows the CRISPRi strategy. **c**, Heat map denoting the log₂-transformed fold change in expression of OPs and ONPs measured after CRISPRi against ONPs (qRT-PCR changes; see Methods). **d**, Top, Hi-C map showing *TERT* and *CLPTM1L* genes in a shared contact domain. Bottom, qRT-PCR results showing the expression of *TERT* and *CLPTM1L* in wild-type and *CLPTM1Lp*-KO 293T cells ($n = 3$ biological replicates). **e**, Cancer mutations in

CLPTM1Lp (each dot represents a mutation ID, with green-coloured mutations predicted to disrupt CTCF motifs (loss), and blue not to disrupt (no loss)). The CTCF motif containing the three mutations is shown at the top. The y axis shows the number of donors for each mutation. Donor IDs, cancer types (abbreviations shown in the Methods) and genomic coordinates are indicated (green). **f**, 293T cells were knocked-in (KI) with these mutations (*KI-CLPTM1Lp*). qRT-PCR results show the RNA expression levels of *CLPTM1L* and *TERT* in wild-type versus *KI-CLPTM1Lp* cells ($n = 3$ biological replicates). **g**, ChIP-qPCR results showing CTCF binding at *CLPTM1Lp* in wild-type versus *KI-CLPTM1Lp* cells ($n = 4$ biological replicates). Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's *t*-test.

level, revealing many significantly mutated ONPs (Supplementary Fig. 6).

To examine whether defects in ONPs might elicit the activation of cancer genes through ERR, we conducted a focused screening. We found that epigenetic inhibition of promoters by dCas9-KRAB/gRNA (CRISPR interference; CRISPRi) recapitulated the ERR effect that was caused by genetic deletion (Fig. 3b). We therefore selected 36 ONPs with cancer mutations or deletions and performed CRISPRi on their promoters to test the effects on nearby oncogenes (Extended Data Fig. 9c, Methods). Effective CRISPRi was achieved for 25 ONPs (Methods). Of these, eight induced a significant increase in the expression of neighbouring oncogenes whereas one led to a significant decrease (Fig. 3c, Supplementary Table 3). For example, inhibition of the *ZCCHC7* and *PVT1* promoters increased the expression of the neighbouring genes *PAX5* and *MYC*, respectively (Extended Data Fig. 9d); the latter is consistent with a previous report¹⁷. Inhibition of the *MTG2* promoter—a recently identified mutation hotspot^{16,18}—led to significant upregulation of a neighbouring cancer gene, *SS18L1*¹⁹ (Fig. 3c, Extended Data Fig. 9d). These results suggest that ERR potentially underlies a subset of noncoding cancer defects and can activate oncogenes.

Deletion or mutation of ONPs activates oncogenes

To better model cancer genetics, we performed deletion or mutation of select ONPs (Supplementary Fig. 5). Three ONP–OP pairs, *PVT1*–*MYC*, *ZCCHC7*–*PAX5* and *CLPTM1L*–*TERT*, were chosen on the basis of reported mutations or deletions and our screening (Methods). Each ONP is located in a shared domain with the respective oncogene (Fig. 3d, Extended Data Fig. 9e, f). Oncoplots of promoter-homed mutations showed that each ONP contains mutations in around 0.5–1% of more than 6,000 tumours (Extended Data Fig. 9g, h, i). Each ONP loops with an adjacent eRNA-expressing enhancer (Extended Data Fig. 10a–c), supporting the possibility that ONP loss might elicit ERR events. Indeed, ONP deletion increased the expression of its respective oncogene by around two- to fivefold (Fig. 3d, Extended Data Fig. 9e, f), and this was accompanied by a gain of interaction between OPs and nearby enhancers (Extended Data Fig. 10d, e).

We found that cancer mutations that overlap CTCF motifs—but not FOXA1, GATA3 or ER α motifs—were common in gene promoters (Extended Data Fig. 10f). Recurrent single-nucleotide mutations in ONPs, as compared to mutations in OPs or random promoters, are more likely to disrupt CTCF motifs (Supplementary Fig. 7b). We identified three cancer mutations in the *CLPTM1L* promoter (*CLPTM1Lp*) overlapping a single CTCF motif, which are predicted to disrupt CTCF binding (Fig. 3e). We generated a knock-in cell line heterozygous for these mutations (Supplementary Fig. 5c), which exhibited an increase of *TERT* but decrease of *CLPTM1L* expression (Fig. 3f) and showed reduced CTCF binding at *CLPTM1Lp* (Fig. 3g). Consistent with this observation in cellular models, data from two individuals with *CLPTM1Lp* mutations showed that these individuals had a level of *TERT* mRNA that was three- to fivefold higher than average (Supplementary Fig. 7c). Together, these results show that specific deletions or single-nucleotide mutations of ONPs can deregulate adjacent oncogenes through ERR, offering mechanistic insights into noncoding mutations and deletions in cancer¹⁶.

ERR functions in some disease-risk loci

We analysed GTEx data to infer potential ERR events in human genetics (Methods, Supplementary Figs. 8–10). We examined GTEx-defined expression quantitative trait loci (eQTLs) in promoters that highly correlate with the expression of both the host gene per se (dubbed gene-CP for cognate promoter) and another gene in the chromosomal neighbourhood (dubbed gene-AP for alternative promoter), in a tissue-specific manner (Extended Data Fig. 11a). This revealed more than 19,000 gene-CP–gene-AP pairs that exhibited opposite trends of allelic expression correlation with eQTLs in gene-CPs (Supplementary Table 4a). We then selected gene-CPs that have one or more additional eQTLs in their 200-kb neighbourhood that happen to overlap annotated enhancers. These criteria identified 872 gene pairs that were potentially undergoing ERR (Supplementary Table 4b), of which 61.4% (535 out of 872) are observed in multiple tissues and 38.6% (337 out of 872) in one tissue (Extended Data Fig. 11b). Each tissue possesses tens to hundreds of potential ERR events (Fig. 4a, Extended Data Fig. 11c).





Fig. 4 | ERR represents an overlooked molecular event that can explain disease susceptibility. **a**, Scatter plot and linear regression showing the numbers of ERR-like events relative to the number of RNA sequencing (RNA-seq) samples in each tissue. The P value is estimated from the t -statistic of the fit. The error band represents the standard error of the regression model. **b**, Box plots showing GTEx normalized effect size (NES) of *NUCKS1* or *RAB7L1* genes in the alternative allele (Alt) relative to the reference allele (Ref), as segregated by the rs823114 SNP. For details of sample numbers, NES, P and m values, see Methods. Cor, cortex; hemi., hemisphere. **c**, SNPs and allele-specific RNA expression (ASE) in *NUCKS1*-*RAB7L1* loci in two lines of iPS cells. Three SNPs at the *NUCKS1* promoter that are associated with Parkinson's



By selecting gene pairs that bear eQTLs in gene-CPs that overlap with risk alleles from genome-wide association studies, we identified 85 potential ERR-based disease-susceptibility events (Supplementary Table 4c). Among these, the *PARK16* locus is a prominent risk locus associated with Parkinson's disease²⁰. *RAB7L1* (also known as *RAB29*) is a key gene in this locus, with a critical role in Parkinson's disease²¹. Notably, three SNPs that are significantly associated with Parkinson's disease risk—rs823114, rs823116 and rs7536483—locate to the promoter of *NUCKS1*, a gene that neighbours *RAB7L1* but which has no reported function in Parkinson's disease (Fig. 4c). In multiple brain tissues, these SNPs are associated with a lower expression of the gene-CP (*NUCKS1*) but a higher expression of the neighbouring disease gene (*RAB7L1*) (Fig. 4b). We identified two clones of human induced pluripotent stem (iPS) cells that contain a heterozygous haplotype of these SNPs. Analysis of allelic expression revealed a lower expression of *NUCKS1* and a higher expression of *RAB7L1* by the alternative allele (Fig. 4c). We differentiated iPS cells to neural progenitor cells (NPCs), a more disease-relevant cell stage (Supplementary Fig. 11), and confirmed the allele-biased expression (Fig. 4e). Allelic 4C-seq in NPCs revealed an E-P interaction between the *NUCKS1* promoter and a putative enhancer in the reference allele, which was lost in the alternative allele; by contrast, the *RAB7L1* promoter loops with this enhancer only in the alternative allele (Fig. 4d). This putative enhancer overlaps H3K27ac signal from the ENCODE project datasets (Fig. 4c), and bears bona fide enhancer activity because its CRISPRi-mediated inhibition decreased the mRNA expression of *NUCKS1* and *RAB7L1* (Extended Data Fig. 11d). Allelic chromatin immunoprecipitation with quantitative PCR (ChIP-qPCR) showed that CTCF binds to a significantly higher level at the reference allele of the *NUCKS1* promoter than at the alternative allele (Fig. 4f, Extended Data Fig. 11e, f). Together, these results indicate that ERR functions in disease-risk loci to control allelic E-P engagement and gene activation, and for at least a subset of loci, this occurs on the basis of altered CTCF binding.

Conclusions

Here, we have delved into the mechanisms that underlie the functional choice of a target gene promoter by a regulatory enhancer, and examined the relevance of these mechanisms to disease. Extensive promoter-disruption experiments revealed that a regulatory enhancer has the potential to contact many or all promoters in its hosting domain¹², but exhibits a 'preference' in choosing one major target promoter to confer the strongest activation. Our data suggest that such promoter preference is determined by inherent features of the promoters and, for at least some loci, by the level of CTCF binding. These results are most compatible with an 'enhancer scanning' model²² as the basis of E-P engagement inside a chromatin domain (Extended Data Fig. 7). Together with the importance of cohesin^{23,24}, we suggest that finer-scale E-P looping is licensed by low-affinity CTCF sites at promoters dynamically interacting with enhancer-bound cohesin to achieve the preferred E-P activation event. Because CTCF binding is observed at a substantial subset of promoters, it apparently represents a mechanism for determining a subset of promoter preference events.

We show that ERR is a clinically important paradigm that underlies genetic or epigenetic alterations near to or within promoters, by which the defects of a primary, non-disease-causing promoter license the activation of alternative, disease-causing gene promoters in a shared chromatin domain. Two notable features of ERR make it applicable to disease: (i) it can cause a striking activation of alternative genes (which as compared to normal conditions are often increased in expression by two- to fivefold and sometimes by around 20-fold); (ii) it potentially has a relatively high prevalence, as suggested by analyses of cancer genomics and GTEx data and by our initial CRISPRi screening. Large-scale promoter perturbation screenings will be required in the future to go beyond informatic inference and fully establish the frequency of ERR events underlying disease gene alterations.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-021-03577-1>.

1. Schoenfelder, S. & Fraser, P. Long-range enhancer–promoter contacts in gene expression control. *Nat. Rev. Genet.* **20**, 437–455 (2019).
2. Hsieh, T. S. et al. Resolving the 3D landscape of transcription-linked mammalian chromatin folding. *Mol. Cell* **78**, 539–553 (2020).
3. Gasperini, M. et al. A genome-wide framework for mapping gene regulation via cellular genetic screens. *Cell* **176**, 377–390 (2019).
4. van Arensbergen, J., van Steensel, B. & Bussemaker, H. J. In search of the determinants of enhancer–promoter interaction specificity. *Trends Cell Biol.* **24**, 695–702 (2014).
5. Li, W. et al. Functional roles of enhancer RNAs for oestrogen-dependent transcriptional activation. *Nature* **498**, 516–520 (2013).
6. Arner, E. et al. Transcribed enhancers lead waves of coordinated transcription in transitioning mammalian cells. *Science* **347**, 1010–1014 (2015).
7. The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57–74 (2012).
8. Rao, S. S. et al. A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**, 1665–1680 (2014).
9. Nora, E. P. et al. Targeted degradation of CTCF decouples local insulation of chromosome domains from genomic compartmentalization. *Cell* **169**, 930–944 (2017).
10. Hnisz, D. et al. Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934–947 (2013).
11. Li, Y. et al. The structural basis for cohesin–CTCF-anchored loops. *Nature* **578**, 472–476 (2020).
12. Lucas, J. S., Zhang, Y., Dudko, O. K. & Murre, C. 3D trajectories adopted by coding and regulatory DNA elements: first-passage times for genomic interactions. *Cell* **158**, 339–352 (2014).
13. Guo, Y. et al. CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter function. *Cell* **162**, 900–910 (2015).
14. Jin, E. H. et al. Association between promoter polymorphisms of TFF1, TFF2, and TFF3 and the risk of gastric and diffuse gastric cancers in a Korean population. *J. Korean Med. Sci.* **30**, 1035–1041 (2015).
15. Weinhold, N., Jacobsen, A., Schultz, N., Sander, C. & Lee, W. Genome-wide analysis of noncoding regulatory mutations in cancer. *Nat. Genet.* **46**, 1160–1165 (2014).
16. Rheinbay, E. et al. Analyses of non-coding somatic drivers in 2,658 cancer whole genomes. *Nature* **578**, 102–111 (2020).
17. Cho, S. W. et al. Promoter of lncRNA gene *PVT1* is a tumor-suppressor DNA boundary element. *Cell* **173**, 1398–1412 (2018).
18. Zhang, W. et al. A global transcriptional network connecting noncoding mutations to changes in tumor gene expression. *Nat. Genet.* **50**, 613–620 (2018).
19. Tate, J. G. et al. COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res.* **47**, D941–D947 (2019).
20. Satake, W. et al. Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease. *Nat. Genet.* **41**, 1303–1307 (2009).
21. Liu, Z. et al. LRRK2 phosphorylates membrane-bound Rabs and is activated by GTP-bound Rab7L1 to promote recruitment to the trans-Golgi network. *Hum. Mol. Genet.* **27**, 385–395 (2018).
22. Blackwood, E. M. & Kadonaga, J. T. Going the distance: a current view of enhancer action. *Science* **281**, 60–63 (1998).
23. Rao, S. S. P. et al. Cohesin loss eliminates all loop domains. *Cell* **171**, 305–320 (2017).
24. Schwarzer, W. et al. Two independent modes of chromatin organization revealed by cohesin removal. *Nature* **551**, 51–56 (2017).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2021

Methods

Data reporting

No statistical methods were used to predetermine sample size. The experiments were not randomized and the investigators were not blinded to allocation during experiments and outcome assessment.

Cell culture and transfection

The protocol for cell culture followed previous work²⁵. In brief, we originally purchased MCF-7 cells from ATCC, which were maintained in DMEM (Gibco 10566) supplemented with 10% FBS (Omega Scientific) in a 5% CO₂ humidified incubator at 37 °C (MCF-7 without stripping). Cells were examined for mycoplasma contamination every 6–12 months. To induce the oestrogen gene transcriptional responses, MCF-7 cells at 60–80% confluency were first hormone-stripped for 3 days in phenol-free DMEM plus 5% charcoal–dextran-treated FBS (Omega Scientific), and they were then treated with 100 nM 17- β -oestradiol (Sigma) for the indicated times (usually 1 h). Control samples were treated with ethanol or ICI 182,780, a high-affinity ER α antagonist. Transfection of siRNAs into MCF-7 cells was performed using lipofectamine 2000 (Life Technologies) following the manufacturer's instructions. For all experiments, two rounds of siRNA transfection (40 nM each time) were performed to achieve higher efficiency. The siRNAs used in this study include: Qiagen negative control siRNA (Qiagen 1027310) and/or Sigma Mission siRNA universal control #2 (SIC002), and siRAD21 (Sigma SASI_Hs02_00341219 and SASI_Hs01_00195799).

CRISPR–Cas9-mediated genome editing

For knockout of genomic regions, two gRNA sequences were designed for each locus with the <http://crispr.mit.edu/> design tool. Point mutations at *CLPTMILP* were achieved using single-stranded oligo DNA nucleotides (ssODNs) as the homology repair template with single-guide RNA (sgRNA) and Cas9 expression. The sgRNA sequences were cloned into a Cas9 vector (pSpCas9(BB)-2A-Puro (PX459) V2.0, 62988) from Addgene (Cambridge), using the BbsI restriction enzyme. The sequences of all sgRNAs and ssODNs used in this study are listed in Supplementary Table 5. Transient transfection of sgRNA–Cas9 plasmids was performed with lipofectamine 2000 to MCF-7 or 293T cells. After a 24-h incubation, the transfected cells were incubated overnight with 2 μ g/ml puromycin. Then cells were plated at clonal density and grown without puromycin until colonies appeared. Individual colonies were picked with a P100 pipette under the observation of a microscope inside a sterile tissue culture hood. Colonies were then moved to individual wells of a 96-well or 48-well plate. Colonies carrying a deletion allele were checked by PCR and confirmed by Sanger sequencing. The sequences of all genotyping primers are listed in Supplementary Table 5.

qPCR

RNA was isolated using TRIzol (Life Technologies), RNeasy Mini Column (Qiagen) or Zymo MiniPrep RNA prep kit (Zymo Research), which was always complemented with DNase treatment. The total RNA was reverse-transcribed using SuperScript III Reverse Transcriptase (Life Technologies) or qScript XLT cDNA SuperMix (QuantaBio) with random hexamer as per the manufacturer's instructions. qPCR was performed in either StepOne Plus or QuantStudio 3 qPCR systems (Applied Biosystems, Thermo Fisher Scientific) using 2X qPCR master mix from Affymetrix (75690) or Applied Biosystems (4472908, SYBR Select). Normalization of expression was done using *GAPDH* or *ACTB* mRNA as internal controls. *P* values were obtained using a two-tailed Student's *t*-test. A list of primers used for qPCR is provided in Supplementary Table 5. For all qRT–PCR and ChIP–qPCR, experiments were performed with at least biological duplicates. Each biological replicate has at least three technical repeats.

ChIP and ChIP–seq

ChIP was performed as previously described²⁵. In brief, cells were cross-linked with 1% formaldehyde at room temperature for 10 min.

Or for some cases, cells were double cross-linked with 1 mM DSG (ProteoChem) for 1 h first and then for 10 minutes by 1% formaldehyde. In both situations, the cross-linking was quenched by addition of 0.125 M glycine for 10 min. ChIP chromatin was fragmented using (Diagenode) Bioruptor300 (20–40 cycles, 30 s on, 30 s off, 4 °C). Subsequently, the soluble chromatin was collected by 16,100g centrifugation, pre-cleared with 10–20 μ l Dynabeads G (Life Technologies), and then incubated with 1–5 μ g of antibodies at 4 °C overnight. The next morning, immunoprecipitated complexes were collected using 30 μ l of Dynabeads protein G (Invitrogen) per reaction. The immune complexes were subjected to washes once with wash buffer I, twice with wash buffer II, once with Tris-EDTA (TE) + 0.1% Triton X-100, and once with TE, and then the beads were incubated at 55 °C for 2 h with proteinase K and de-cross-linked at 65 °C overnight. The final ChIP DNA was extracted and purified using QIAquick spin columns (Qiagen). For ChIP–seq, the extracted DNA was ligated to specific adaptors for Illumina's HiSeq system using the KAPA Hyper Prep Kit (Kapa Biosystems).

High-throughput sequencing

For all ChIP–seq and GRO–seq, the extracted DNA libraries were sequenced with the Illumina HiSeq 2500 or HiSeq 4000 system according to the manufacturer's instructions. The first 50 bp for each sequence tag returned by the Illumina Pipeline was aligned to the human genome (hg19) assembly using Bowtie2, and only reads with MAPQ $\geq$ 10 were selected by SAMtools for further analysis. The data were visualized by preparing custom tracks on the University of California, Santa Cruz (UCSC) genome browser using HOMER²⁶ (<http://homer.ucsd.edu/homer/>). The total number of mappable reads was normalized to 10⁷ for each experiment presented in this study. The read numbers used for generating bedgraph files are provided in Supplementary Table 6.

4C-seq and analyses

The 4C-seq experiments were conducted following a published protocol²⁷ with modification. In brief, 10 million cells were cross-linked with 1% formaldehyde for 10 min and nuclei were extracted. Nuclei were resuspended in restriction enzyme buffer and incubated with 0.3% SDS for 1 h at 37 °C and further incubated with 2% Triton X-100 for 1 h; then 400 U of DpnII restriction enzyme was added and nuclei were incubated overnight. Restriction enzyme was heat-inactivated at 65 °C for 20 min. Ligation of DNA regions in close physical proximity was performed using 1,000 U of T4 DNA ligase (NEB) for overnight. After de-cross-linking, the second digestion and ligation were performed using restriction enzyme NlaIII and T4 DNA ligase. The 4C-seq libraries were amplified using PCR with the first primer designed on each viewpoint and the second primer designed beside the NlaIII site. Both primers contained Illumina sequencing adaptors and barcode (Supplementary Table 5). The 4C libraries were sequenced on the Illumina HiSeq 2500 using single-read 100-cycle runs.

Analysis of 4C-seq data was done using an existing 4C pipeline²⁸. Valid reads of genomic regions were generated by clipping out the primer sequences in the raw reads. These clipped reads with unique 3' ends were aligned to hg19 human genomic coordinates. All the samples were quality checked according to the following: ratio of the read number of *cis*-interactions versus the read number of total-interaction is larger than 40%. For data visualization, a 5 kb window size was chosen to compute the trend curve and the grey band on top of each 4C heat map displays the 20–80% for the windows. Median contact intensities were depicted as colour-coded multi-scaled heat maps, ranging from a 2-kb sliding window at the top of the heat map to a 50-kb sliding window at the bottom of the heat map. At *NUCKS1* promoter and *RAB7LI* promoter viewpoints, rs823116 and rs708725 were used for detection of allele-specific interaction, respectively.

Replicates of 4C-seq data were processed following a previous method for statistical analysis²⁹. In brief, 4C-seq reads were first trimmed by keeping the sequences after restriction enzyme cutting

Article

site 1 and before restriction enzyme cutting site 2. Trimmed reads were mapped to the hg19 reference genome with Bowtie2. Mapped reads were counted for each restriction fragment end. For each fragment end, the read counts were further normalized by scaling the number of total reads mapped to the viewpoint chromosome. After excluding the reads mapped to the top 2 fragments, the normalization will make the total sum of normalized reads 1 million (normalization factor). For plots for which quantitation bar plots were included, normalized read counts of all valid fragment ends overlapping *NUCKS1e*, *TFF1p*, *KCNK5p* and *P2RY2p* were summed up and compared by Student's *t*-test.

GRO-seq

GRO-seq experiments were performed as previously reported²⁵. In brief, around 10–20 million MCF-7 cells were washed 3 times with cold PBS and then sequentially swelled in swelling buffer (10 mM Tris-Cl pH 7.5, 2 mM MgCl₂, 3 mM CaCl₂) for 5 min on ice, collected, and lysed in lysis buffer (swelling buffer plus 0.5% NP-40 and 10% glycerol). The resultant nuclei were washed one more time with 10 ml lysis buffer and finally resuspended in 100 µl of freezing buffer (50 mM Tris-Cl pH 8.3, 40% glycerol, 5 mM MgCl₂, 0.1 mM EDTA). For the run-on assay, resuspended nuclei were mixed with an equal volume of reaction buffer (10 mM Tris-Cl pH 8.0, 5 mM MgCl₂, 1 mM DTT, 300 mM KCl, 20 units of SUPERase-IN, 1% sarkosyl, 500 µM ATP, GTP and Br-UTP, 2 µM CTP) and incubated for 5 min at 30 °C. The resultant nuclear-run-on RNA (NRO-RNA) was then extracted with TRIzol LS reagent (Life Technologies) following the manufacturer's instructions. NRO-RNA was fragmented to around 300–500 nt by alkaline base hydrolysis on ice and followed by treatment with DNase I and Antarctic phosphatase. The fragmented Br-UTP-labelled nascent RNA was then immune-precipitated with anti-BrdU agarose beads (Sc32323ac, Santa Cruz Biotechnology) in binding buffer (0.5× SSPE, 1 mM EDTA, 0.05% Tween) for 3 h at 4 °C with rotation. Subsequently, T4 PNK was used to repair the end of the immune-precipitated BrU-NRO-RNA, at 37 °C for 1 h. The RNA was extracted and precipitated using acidic phenol–chloroform.

cDNA synthesis was performed as per a published method³⁰ with a few modifications. The RNA fragments were subjected to poly-A tailing reaction by poly-A polymerase (NEB) for 30 min at 37 °C. Subsequently, reverse transcription was performed using oNTI223 primer and superscript III RT kit (Life Technologies). The cDNA products were separated on a 10% polyacrylamide TBE-urea gel and only those migrating between around 100–500 bp were excised and recovered by gel extraction. After that, the first-strand cDNA was circularized by CircLigase (Epicentre) and re-linearized by APE1 (NEB). Re-linearized single-strand cDNA (sscDNA) was separated by a 10% polyacrylamide TBE gel as described above and the product of required size was excised (around 170–400 bp) for gel extraction. Finally, the sscDNA template was amplified by PCR (usually between 10–14 PCR cycles) using the Phusion High-Fidelity enzyme (NEB) according to the manufacturer's instructions. The examples of oligonucleotide primers oNTI200 (with 4 barcode choices) and oNTI201 were used to generate DNA for deep sequencing (sequences are listed below from 5' to 3' direction).

oNTI223-TruSeq: /5Phos/GATCGTCGGACTGTAGAAGCTCT; CAGACGTGTGCTCTTCCGATCTTTTTTTTTTTTTTTTTTTVN (";" = abasic dSpacer furan; VN = degenerate nucleotides). oNTI200-TruSeqID1: CAAGCAGAAGACGGCATACGAGTGTGACTGGAGTTCAGACGTGCTCTTCCGATC; oNTI200-TruSeqID2: CAAGCAGAAGACGGCATACGAGTGTGACTGGAGTTCAGACGTGCTCTTCCGATC; oNTI200-TruSeqID3: CAAGCAGAAGACGGCATACGAGTGTGACTGGAGTTCAGACGTGCTCTTCCGATC; oNTI200-TruSeqID4: CAAGCAGAAGACGGCATACGAGTGTGACTGGAGTTCAGACGTGCTCTTCCGATC; oNTI201: AATGATACGGCGACCACCGAGGTTCTACAGTCCGACG. Illumina small RNA-seq primer (for first read): CGACAGTTTCAGATTCTACAGTCCGACGATC; index reading primer (for index read): GATCGGAAGAGCACAGTCTGAACCTCCAGTCAC.

GRO-seq analysis

Data were aligned to human genome hg19 by Bowtie2. For gene transcription, the GRO-seq reads from the entire gene bodies of RefSeq genes were included for analysis. For eRNA calculation, we first selected all p300-bound peaks excluding those in gene promoters and transcription termination sites (TTSs) (± 2 kb of TSSs, and $+3$ kb of TTS) as putative enhancers and counted the GRO-seq reads in the ± 2 kb region surrounding the p300 peak summits (called by MACS2). For intergenic enhancers, we counted the sense and anti-sense strands of GRO-seq signals separately, whereas for the intragenic enhancers, only the strand opposite to the coding gene was considered. For counting differential gene or eRNA expression, raw read counts from GRO-seq were generated using HOMER (analyzeRepeats.pl -raw), and subjected to analysis by EdgeR to find differentially expressed (DE) genes or eRNAs (false discovery rate (FDR) < 0.05 , fold change ≥ 1.5 or ≤ 0.667). The enhancers with upregulated eRNAs were selected and the tag counts of epigenomic marks (for example, H3K4me3 or H3K4me1) on these enhancers were generated by HOMER²⁶ and then plotted as profile plots using the Bioconductor package in R.

CRISPRi and tethering

To achieve efficient silencing at each target locus, we used two sgRNAs in one plasmid system. gRNAs were designed for each targeted locus using the Alt-R Custom Cas9 crRNA Design Tool (<https://eu.idtdna.com/>) and cloned into an in-house lentiviral vector pLKO.1-U6-2sgRNA-ccdB-E F1a-Puromycin (which can express two gRNAs) in one vector. Lentiviral gRNAs or Lenti-dCas9-KRAB-blast plasmids (89567, Addgene) were co-transfected with packaging plasmids (psPAX2 and pMD2.G) into HEK-293-T cells using lipofectamine 2000. Culture medium containing lentivirus for gRNA and dCas9-KRAB was collected and added to the desired cells with 8 µg/ml polybrene. Next day, the medium was changed with medium containing appropriate antibiotics. After 48 h of drug selection, cells were collected for further experiments. The list of oncogenes was selected mostly on the basis of the COSMIC Cancer Gene Census (CGC)¹⁹ Tier 1 genes, preferentially those with annotation to bear cancer amplification. We selected these that contain cancer mutations in ONPs based on our own analysis (Extended Data Fig. 9c, Supplementary Fig. 6). Some ONP-OP pairs—for example, *ZCCHC7-PAX3*³¹, *CLPTMIL-TERT*¹⁵, *PVT1-MYC*¹⁷, *MYL12A-TGIF1* or *MYL12B-TGIF1*, *ID1-PIM1*, *HMI3-PIM1*³² and *MTG2-SS18L*^{16,18}—were also included on the basis of literature reports of their potential importance in cancer. In addition, ERR predicts that the functional loss of ONPs will upregulate OPs, which requires ONPs to be transcribed. Therefore, in our selection of 36 pairs, we require both ONP-OP genes to display detectable GRO-seq signals in MCF-7 cells, particularly for the ONP genes. For the focused CRISPRi screening, all the gene-expression changes and *P* values of ONPs and OPs are shown in Supplementary Table 3. We only considered that CRISPRi for ONP genes worked if a fold reduction of ≤ 0.667 and a *P* value < 0.05 were observed; subsequently, we counted the numbers of up- or downregulated OP genes by expression fold change of ≥ 1.5 or ≤ 0.667 as well as $P < 0.05$. *P* values here denote two-tailed Student's *t*-tests. For Fig. 3c, we plotted the log₂-transformed fold change of 25 ONP-OP gene pairs with a red–blue gradient.

For CTCF tethering, we used the SunTag system³³. We generated dCas9-10xGCN4 and scFv-CTCF constructs and expressed them together with specific gRNAs targeting the TFF3p (Extended Data Fig. 6e, f). MCF7 cells were transduced with lentivirus expressing dCas9-10xGCN4-T2A-blasticidin and scFV-CTCF_wildtype-T2A-hygromycin or scFV-CTCF_Y226A_F228A-T2A-hygromycin. The cells were selected with hygromycin-B (200 µg/ml) and blasticidin (10 µg/ml) for 7 days. dCas9-10xGCN4 and scFV-CTCF (wild-type or Y226A_F228A) stably expressing cell lines were then transduced with lentivirus expressing target sgRNAs, and the sgRNA-positive cells were further selected with puromycin (2 µg/ml) for 2 days in the presence of hygromycin-B and blasticidin.

Generation and culture of NPCs

We identified two clones of human iPS cells that contain a heterozygous haplotype of Parkinson's disease-risk SNPs, rs823114³⁴, rs823116³⁵ and rs7536483³⁶, in the promoter of *NUCKS1*. We generated monolayers of NPCs from identified human iPS cell cultures (iPSCORE_19_1_iPSC_C4_P13 and iPSCORE_2_11_iPSC_C2_P12 human iPS cell lines) using the serum-free medium kit for highly efficient SMAD inhibition-mediated neural induction and following the manufacturer's recommendations (STEMdiffTM SMADi Neural Induction Kit, 08581, from STEMCELL Technologies; which contains STEMdiffTM Neural Induction Medium and STEMdiffTM SMADi Neural Induction Supplement). SMAD inhibitors were intended to inhibit the differentiation to a non-CNS-type of cells, according to STEMCELL Technologies. In brief, we thawed a vial of frozen human iPS cells in a well of a 6 well-plate (passage 1) pre-coated with MatrigelR (Corning, 354277). The human iPS cell medium was mTeSR1TM from STEMCELL Technologies supplemented with 20 mM penicillin–streptomycin (Gibco, 15140122). We used 1:1,000 Y-27632 ROCK Inhibitor (Abcam-ab 120129) for 24 h to increase the survival of single human iPS cells (not used for daily medium changes). We passaged human iPS cells at a 1:3 dilution twice after 4 days of culturing every time (passages 2 and 3). For NSC induction, we plated 2 million human iPS cells in a well of a 6-well plate (from passage 3), detaching cells using Versene and disrupting at the single-cell level. Cell counts were performed using Trypan blue staining and the Countess II Automated Cell Counter instrument from Life Technologies. We started NSC induction with the STEMdiff SMADi Neural Induction Kit. Six days later, we passaged cells at a 1:6 dilution using Accutase (Innovative Cell Technologies, AT104) (passage 5). Six days later, NSCs were passaged again (passage 6) at a 1:6 dilution in 10-cm plates. NSC medium was changed daily in all cases. Five days later, NSCs were collected for further experiments. In total, the period of NSC induction was 17 days. We confirmed the NSC identity by RNA-seq (Supplementary Fig. 11). As expected, we did not observe expression of pluripotency markers *NANOG* and *POU5F1* (OCT4), but markers of (neural) stem cell identity, *NES* (nestin), *MKI67* (Ki67), *LIN28B* and *SOX2* were found. We observed robust expression of neuronal-lineage markers *TUBB3* (TUJ1), *ENO2* (NSE) and *MAP2*, but no or relatively low expression of specialized genes of neuronal identity (such as *MAPT*, *SYP* or *TH*) or glial differentiation (such as *GFAP* and *S100B*). Together, these results corroborate that our cells have NPC identity.

iPS cell and whole-genome sequencing data from two individuals

The two individuals (iPSCORE_2_11 and iPSCORE_19_1) were recruited as part of the iPSCORE project³⁷. The recruitment of these individuals was approved by the Institutional Review Boards of the University of California, San Diego and The Salk Institute (project no. 110776ZF). As previously described³⁷, we generated whole-genome sequences from DNA isolated from blood on the HiSeqX (Illumina; 150-bp paired end). Whole-genome sequencing data are available at the National Institutes of Health (NIH) dbGaP database (phs001325).

iPS cell generation

Cultures of primary dermal fibroblast cells were infected with the Cyto-tune Sendai virus (Life Technologies) per the manufacturer's protocol. The Sendai-infected cells were maintained with 10% FBS–DMEM (Invitrogen) for days 4–7 until the cells recovered and repopulated the well. Emerging iPS cell colonies were manually picked after day 21 and maintained on Matrigel (BD Corning) with mTeSR1 medium (STEMCELL Technologies) as previously described³⁸. Multiple independently established iPS cell lines (that is, referred to as clones) were derived from each individual. Sendai virus clearance typically occurred at or before passage (P) 9, and was not detected in the iPS cell lines at the P12 stage of cryopreservation. RNA was collected at P12 for iPSCORE_2_11 (iPSC.ID = iPSCORE_2_11_iPSC_C2_P12) and at P13 for iPSCORE_19_1 (iPSC

ID = iPSCORE_19_1_iPSC_C4_P13). Both iPS cell samples were pluripotent³⁹. iPS cell lines are publicly available through the NHLBI-contracted biorepository at the WiCell Research Institute.

iPS cell RNA library preparation and sequencing

iPS cell RNA-seq data were generated as previously described³⁷. In brief, total RNA was extracted from the iPS cell lines using AllPrep RNasy Blood & Tissue Kit (Qiagen) and the quality was assessed based on RNA integrity number (RIN) using an Agilent Bioanalyzer. Libraries were prepared using the Illumina TruSeq stranded mRNA kits and sequenced using an Illumina HiSeq2500 (around 11 samples per lane). Samples were sequenced to an average of around 22 million read pairs. RNA-seq data are available through dbGaP (phs000924).

Allele-specific expression

iPS cell allele-specific expression (ASE) was calculated as previously described³⁷ (shown in Fig. 4c). RNA-seq reads were aligned to the hg19 genome using STAR⁴⁰ and sorted using Sambamba⁴¹. Biobambam2 bam-markduplicates was used to mark duplicate reads. Uniquely mapped reads that were not marked as duplicates were tested for mapping bias using WASP⁴². GATK ASEReadCounter was used to calculate the coverage of heterozygous variants on Gencode V.19 exons⁴³. All heterozygous variants with coverage ≥ 8 , reference allele frequency between 2% and 98%, located in uniquely mappable regions according to the wgEncodeCrgMapabilityAlign100-mer track and >10 bp away from other variants, were tested for ASE using MBASED⁴⁴. NPC ASE and allele-specific CTCF binding were measured by the rhAmp SNP Genotyping System (IDT) according to the manufacturer's instructions, and were based on qPCR. rs111265946 and rs823137 were used for detection of *NUCKS1* and *RAB7L1* ASE, respectively. In brief, the total RNA was reverse-transcribed using SuperScript III Reverse Transcriptase (Life Technologies). qPCR was performed with rhAmp Genotyping Master Mix and rhAmp Reporter Mix (IDT). FAM and VIC were assigned to the reference allele and alternative allele, respectively. Expression was normalized to qPCR of genomic DNA. rs823114 was used for detection of allele-specific CTCF binding at the *NUCKS1* promoter (Fig. 4f). ChIP was performed as described above and ChIP–qPCR was performed with rhAmp Genotyping Master Mix and rhAmp Reporter Mix (IDT). FAM and VIC were assigned as the reference allele and alternative allele, respectively. The relative quantities of ChIP samples were normalized by individual input DNA samples.

Other bioinformatic analyses

Many published datasets are used for our analyses (Supplementary Table 6). A description of the raw data generated in this paper is also included as Supplementary Table 6.

For Fig. 2a, CTCF ChIP–seq peaks were divided into three groups by their locations. Enhancer CTCF were those peaks overlapped with p300-bound enhancers in MCF-7, and promoter CTCF were those overlapped with promoters (RefSeq gene TSSs ± 1 kb); domain boundaries were defined by all the CTCF peaks in the ± 3 kb regions near the TAD boundaries. For each group, HOMER²⁶ was used to calculate the normalized binding profile. For Fig. 2b, after sorting the transcriptional signals of genes identified by GRO-seq, the highest, middle and lowest 1,000 transcribed genes were chosen to find whether there are any CTCF peaks in the promoter (± 1 kb of TSSs). The peak file of CTCF ChIP–seq was downloaded from ENCODE (ENCFF586KIH). A two-sided Fisher's exact test was performed to compare the enrichment of peaks in the promoters for the three groups. For Fig. 4a, we fitted a linear model to estimate the relationship between tissue sample size and the number of ERRs found in that tissue type. Default R functions were used for computing the linear model and the *P* value for the fit (lm), as well as for the calculation of the Pearson's correlation coefficient (*r*). The shaded region around the regression line represents the standard error of the fit. The *P* value for the linear model in R (abbreviated as $\text{Pr}(>|t|)$) is

estimated from the t -statistic of the fit. For Fig. 4b showing the normalized effect size (NES) of GTEx eQTL SNPs, the definitions of NES, P value and m value can be found in the GTEx portal (<https://www.gtexportal.org/>), V8. We quote the definition here. For m value, it denotes the posterior probability that an eQTL effect exists in each tissue tested in the cross-tissue meta-analysis; the m value ranges between 0 and 1. For P value, it was generated by a t -test that compares the observed NES from single-tissue eQTL analysis to a null NES of 0. The NES was used to denote the slope of the linear regression of normalized expression data versus the three genotype categories using single-tissue eQTL analysis, representing eQTL effect size. The normalized expression values are based on quantile normalization within each tissue, followed by inverse quantile normalization for each gene across samples.

For Extended Data Fig. 6a–d, super-enhancers and typical enhancers were called by HOMER²⁶ using p300 ChIP-seq datasets in each cell type (ENCFF000QPA for MCF-7, ENCFF258NYC and ENCFF728IVZ for GM12878). For each promoter, the nearest enhancer was identified by BEDtools. For statistics, a two-sided Fisher's exact test was performed in R. For Extended Data Fig. 8a, the enhancers with upregulated eRNAs, as well as promoters from genes downregulated after siRAD21 treatment (as shown in Extended Data Fig. 3) were used to check whether these promoters and enhancers form reciprocal loops in MCF-7 ChIA-PET datasets (GSE39495). Each anchor of the ChIA-PET tag was extended by 5 kb on each side. If an enhancer or a promoter overlaps with the two anchors of a ChIA-PET tag, this pair of enhancer and promoter would be considered a 'looped pair'. A two-sided Fisher's exact test was performed to compare the enrichment of the loops formed by these E-P pairs in MCF-7 cells versus K562 cells. For Extended Data Fig. 8b, c, motif sites for ESRI (MA0112.3) and FOXA1 (MA0148.3) were identified by FIMO. FOXA1 ChIP-seq peaks were obtained from ENCODE (ENCFF596OJV). The enhancer or promoter with both CTCF peaks and motifs were then considered as CTCF-associated enhancer or promoter; the same was done for ESRI and FOXA1.

For Extended Data Figs. 9a–c, g–i, 10f, whole-genome cancer mutation datasets were obtained from the ICGC (release 28). In total, there were 58,402,698 mutations from 2,731 patients as generated from whole-genome sequencing. Only those cancer types with more than 100 donors and more than 1,000 total somatic mutations were used for plotting Extended Data Fig. 9a. For Extended Data Fig. 10f, motifs of CTCF (JASPAR ID MA0139.1), FOXA1 (MA0148.3), GATA3 (MA0037.2) and ESRI (MA0112.3) were identified in the whole genome using FIMO⁴⁵ to find the motif positions. Random genomic sequences with the exact same length as the CTCF motif were generated using BEDTools⁴⁶, and defined as 'random control motifs'. If the distance from a cancer mutation to its closest CTCF motif centre was smaller than 20 bp, the mutation would then be counted as a mutation associated with CTCF motif (that is, CTCF mutations). The mutations in 'random control motif' or the FOXA1, ESRI and GATA3 motifs were defined in the same way. The count of CTCF-motif-disrupting mutations (for example, Fig. 3e) requires that no FIMO CTCF motif can be identified from the mutation-containing sequence anymore. Cancer-type abbreviations can be found in the ICGC portal (<http://icgc.org/>), and include (for example, in Fig. 3e, Extended Data Fig. 9a, c): CLLE-ES, chronic lymphocytic leukaemia (Spain); MALY-DE, malignant lymphoma (Germany); PBCA-DE, paediatric brain cancer (Germany); PBCA-US, paediatric brain cancer (USA); PEME-CA, paediatric medulloblastoma (Canada); BRCA-EU, ER⁺ and HER2⁺ breast cancer (European Union or UK); ESAD-UK, oesophageal adenocarcinoma (UK); LICA-CN, liver cancer (China); LIRI-JP, liver cancer (RIKEN, Japan); PACA-AU, pancreatic cancer (Australia); PACA-CA, pancreatic cancer (Canada); EOPC-DE, early-onset prostate cancer (Germany); PARD-CA, prostate cancer (Canada); PRAD-UK, prostate cancer (UK); MELA-AU, melanoma (Australia); SKCA-BR, skin adenocarcinoma (Brazil).

GTEx data processing

For a detailed description of the method used for GTEx data analysis, see Supplementary Figs. 8–10 and associated notes. The GTEx data used

for the analyses described in this manuscript were obtained from the dbGap accession phs000424.v7.p2.c1 on 12 January 2017. In brief, the search for ERR events in the general human population was conducted using data from the Genotype-Tissue Expression (GTEx) project⁴⁷ V7 release. We searched for *cis*-eQTLs in promoter regions, 2 kb upstream and 1 kb downstream of TSSs. We refer to these genes that host *cis*-eQTLs in their promoters as 'gene-CP' (that is, cognate promoter), and these eQTLs are referred to as P-eQTLs (Extended Data Fig. 11a). We then select the subset of gene-CP promoters for which the P-eQTLs that they contain also act as eQTLs for a distal gene in their chromosomal neighbourhood (± 200 kb) in the same tissue type (this distal gene is referred to as 'gene-AP' for alternative promoter). Although we identified unique gene-CP–gene-AP pairs, we counted the events based on gene names rather than the numbers of P-eQTLs. For example, if 'Gene A' and 'Gene B' are regarded as a unique gene-CP–gene-AP pair, then Gene A/Gene B is counted only once. Next, we identified GTEx *cis*-eQTLs that overlapped with an exhaustive set of experimentally identified enhancer regions annotated by several consortia (the ENCODE¹⁰, FANTOM5³¹, and Roadmap Epigenomics³² projects). The associations with traits and diseases were identified using genome-wide association studies (GWASs) in the NHGRI-EBI GWAS Catalog⁴⁸ and GWASdb v2⁴⁹.

Statistics

In most of the paper, particularly for qRT-PCR results, two-tailed Student's t -tests were used to calculate statistical significance unless otherwise specified. The distribution of sample measurements was considered to follow a normal distribution and variance was assessed as equal. For qRT-PCR or ChIP-qPCR, at least three biological replicates were conducted. For each biological repeat, three technical repeats were performed. Most of the results shown represent mean $\pm$ s.d., with asterisks indicating P values; * P < 0.05, ** P < 0.01, *** P < 0.001 (two tailed Student's t -tests).

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this paper.

Data availability

The datasets generated by the current study are available in the GEO (GSE115604) and the Sequence Read Archive (PRJNA412021). For analyses of cancer mutations and GTEx data, codes are deposited in GitHub (<https://github.com/wblilab-uth/ERR-project> and <https://github.com/zdz-lab/ERR>). Other analyses of ChIP-seq peaks, 4C-seq or gene expression in the current study used standard bioinformatics tools and codes, which are available upon request. Source data are provided with this paper.

- Li, W. et al. Condensin I and II complexes license full estrogen receptor α -dependent enhancer activation. *Mol. Cell* **59**, 188–202 (2015).
- Heinz, S. et al. Simple combinations of lineage-determining transcription factors prime *cis*-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589 (2010).
- Stadhouders, R. et al. Multiplexed chromosome conformation capture sequencing for rapid genome-scale high-resolution detection of long-range chromatin interactions. *Nat. Protocols* **8**, 509–524 (2013).
- van de Werken, H. J. et al. Robust 4C-seq data analysis to screen for regulatory DNA interactions. *Nat. Methods* **9**, 969–972 (2012).
- Krijger, P. H. L., Geeven, G., Bianchi, V., Hilvering, C. R. E. & de Laat, W. 4C-seq from beginning to end: a detailed protocol for sample preparation and data analysis. *Methods* **170**, 17–32 (2020).
- Ingolia, N. T., Ghaemmaghami, S., Newman, J. R. & Weissman, J. S. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* **324**, 218–223 (2009).
- Hayward, N. K. et al. Whole-genome landscapes of major melanoma subtypes. *Nature* **545**, 175–180 (2017).
- Schuijers, J. et al. Transcriptional dysregulation of MYC reveals common enhancer-docking mechanism. *Cell Rep.* **23**, 349–360 (2018).
- Tanenbaum, M. E., Gilbert, L. A., Qi, L. S., Weissman, J. S. & Vale, R. D. A protein-tagging system for signal amplification in gene expression and fluorescence imaging. *Cell* **159**, 635–646 (2014).

34. MacLeod, D. A. et al. RAB7L1 interacts with LRRK2 to modify intraneuronal protein sorting and Parkinson's disease risk. *Neuron* **77**, 425–439 (2013).
35. Latourelle, J. C. et al. Large-scale identification of clinical and genetic predictors of motor progression in patients with newly diagnosed Parkinson's disease: a longitudinal cohort study and validation. *Lancet Neurol.* **16**, 908–916 (2017).
36. Pierce, S. E., Tyson, T., Booms, A., Prah, J. & Coetzee, G. A. Parkinson's disease genetic risk in a midbrain neuronal cell line. *Neurobiol. Dis.* **114**, 53–64 (2018).
37. DeBoever, C. et al. Large-scale profiling reveals the influence of genetic variation on gene expression in human induced pluripotent stem cells. *Cell Stem Cell* **20**, 533–546 (2017).
38. Panopoulos, A. D. et al. The metabolome of induced pluripotent stem cells reveals metabolic changes occurring in somatic cell reprogramming. *Cell Res.* **22**, 168–177 (2012).
39. Panopoulos, A. D. et al. iPSCORE: a resource of 222 iPSC lines enabling functional characterization of genetic variation across a variety of cell types. *Stem Cell Rep.* **8**, 1086–1100 (2017).
40. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
41. Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba: fast processing of NGS alignment formats. *Bioinformatics* **31**, 2032–2034 (2015).
42. van de Geijn, B., McVicker, G., Gilad, Y. & Pritchard, J. K. WASP: allele-specific software for robust molecular quantitative trait locus discovery. *Nat. Methods* **12**, 1061–1063 (2015).
43. Van der Auwera, G. A. et al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1–11.10.33 (2013).
44. Mayba, O. et al. MBASED: allele-specific expression detection in cancer tissues and cell lines. *Genome Biol.* **15**, 405 (2014).
45. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017–1018 (2011).
46. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).
47. Battle, A., Brown, C. D., Engelhardt, B. E. & Montgomery, S. B. Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).
48. MacArthur, J. et al. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res.* **45**, D896–D901 (2017).
49. Li, M. J. et al. GWASdb v2: an update database for human genetic variants identified by genome-wide association studies. *Nucleic Acids Res.* **44**, D869–D876 (2016).
50. Nair, S. J. et al. Phase separation of ligand-activated enhancers licenses cooperative chromosomal enhancer assembly. *Nat. Struct. Mol. Biol.* **26**, 193–203 (2019).
51. Krietenstein, N. et al. Ultrastructural details of mammalian chromosome architecture. *Mol. Cell* **78**, 554–565 (2020).
52. Fudenberg, G. et al. Formation of chromosomal domains by loop extrusion. *Cell Rep.* **15**, 2038–2049 (2016).
53. Hatzis, P. & Talianidis, I. Dynamics of enhancer–promoter communication during differentiation-induced gene activation. *Mol. Cell* **10**, 1467–1477 (2002).

Acknowledgements This work is supported by the Howard Hughes Medical Institute and NIH grants (DK018477, DK039949, HL150521 and NS093066) to M.G.R.; by NIH grants (K22CA204468, R21GM132778, R01GM136922 and '4D nucleome' U01HL156059), a University of Texas Rising STARS Award, the Welch Foundation (AU-2000-20190330), the Gulf Coast Consortium John S. Dunn Foundation and a Cancer Prevention and Research Institute of Texas (CPRIT) Award (RR160083) to W.L.; by NIH grants (HG008118, HL107442, DK105541, DK112155 and NSF-CMMI division award 1728497) to K.A.F.; and by NIH NHGRI grants (R01HG008153 and R01HG008153-S1) to Z.D.Z. W.L. is a CPRIT Scholar in Cancer Research. M.G.R. is an investigator with the Howard Hughes Medical Institute. We thank J. Hightower for assistance with figure preparation, and we acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin (<http://www.tacc.utexas.edu>) for providing high-performance computing resources for part of this work.

Author contributions W.L., S.O. and M.G.R. conceived the project. S.O. and W.L. conducted most experiments, with contributions from X.Z., J.-H.L., S.J.N., F.Y. and K.O. J.S., W.L. and S.O. conducted most of the bioinformatic analyses of this paper, except for the GTEx related analyses, which were done by J.M. and Z.D.Z. M.D. and K.A.F. generated and analysed the RNA-seq data for allelic gene expression in human iPS cells. The 4C-seq data were analysed by F.X., R.W. and Q.M. I.G.-B. contributed to the iPS-cell-derived NPC culture and RNA-seq. W.L., M.G.R. and S.O. wrote the manuscript, with input from all authors.

Competing interests The authors declare no competing interests.


Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41586-021-03577-1>.

Correspondence and requests for materials should be addressed to W.L. or M.G.R.

Peer review information *Nature* thanks Peter Scacheri and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.


Reprints and permissions information is available at <http://www.nature.com/reprints>.



Extended Data Fig. 1 | See next page for caption.

Extended Data Fig. 1 | Disrupted enhancer–promoter looping alters eRNA and mRNA transcription. **a**, qRT–PCR results showing the temporal expression kinetics of eRNAs and mRNAs at two loci after treatment with 17- β -oestradiol (E_2). mRNAs measured by intronic primers ($n = 3$; biological replicates). Shaded error band represents mean $\pm$ s.d. **b**, 4C-seq heat maps using *TFF1e*, *P2RY2e* or *KCNK5e* as the viewpoints showing the effects of *RAD21* siRNA depletion on the E-P looping events; yellow and pink highlights depict enhancers and promoters, respectively. The quantified chromosome contact frequency between enhancer and promoter from two replicates of 4C-seq is shown in the bar graph (bottom). **c**, Expression of *TFF1* mRNA (*TFF1m*) and *TFF1* enhancer RNA (*TFF1e*) in wild-type ($n = 6$) and *TFF1p-KO* MCF-7 isogenic cell clones ($n = 14$) was checked by qRT–PCR. The *FOXCI* gene that locates on another chromosome was examined as a control. The WT5 and KO6 lines were


used for many subsequent experiments (for example, 4C-seq, GRO-seq and ChIP). Each bar represents data from an independent cell clone and was made from $n = 3$ data points of technical replicates. The data presented here represent three biological replicates. **d**, qRT–PCR results showing the expression of *TFF1* eRNA and mRNA in wild-type MCF-7 cells versus *TFF1p-KO* cells with two different gRNA pairs (*TFF1p-KO_gRNA1/2* and *TFF1p-KO_gRNA3/4*) ($n = 3$ data points of technical replicates; representative of three independent experiments). **e**, ChIP–qPCR data indicating the binding of RNA Pol II and p300 at the *TFF1* enhancer or promoter in wild-type compared to *TFF1p-KO* cells ($n = 3$ data points of technical replicates; representative of two biological replicates); their binding at an enhancer region near the *NR1P1* gene (that is, *NR1P1e*) is shown as a control. Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's *t*-test.



Extended Data Fig. 2 | See next page for caption.

Extended Data Fig. 2 | Promoter deletion affects gene and eRNA transcription in its chromosomal neighbourhood at the *P2RY2*, *KCNK5* and *PGR* loci. **a, b**, qRT-PCR showing the mRNA and eRNA expression of *P2RY2* (**a**) and the expression of neighbouring genes (*FCHSD2* and *P2RY6*) (**b**) in wild-type ($n = 2$) versus *P2RY2* promoter KO (*P2RY2p-KO*) ($n = 2$) independent isogenic clones of MCF-7 cells. Each bar represents an independent cell clone ($n = 3$ data points of technical replicates; representative of two independent experiments). **c**, Top, Hi-C contact matrix and ChIP-seq tracks of CTCF and p300 showing the topology of the chromosomal neighbourhood of *P2RY2* locus. Bottom, UCSC browser screenshots showing the binding of CTCF, cohesin subunit (that is, RAD21) and ER α at *P2RY2* and surrounding gene promoters. **d, e**, qRT-PCR showing the mRNA and eRNA expression of *KCNK5* (**d**) and the expression of neighbouring genes (*KCNK17* and *SAYS1*) (**e**) in wild-type ($n = 2$) versus *KCNK5* promoter KO (*KCNK5p-KO*) ($n = 2$) independent isogenic cell clones of MCF-7 cells. Each bar represents an independent cell

clone ($n = 3$ data points of technical replicates; representative of two independent experiments). **f**, Top, Hi-C contact matrix and ChIP-seq tracks of CTCF and p300 showing the topology of the chromosomal neighbourhood of the *KCNK5* locus. Bottom, UCSC browser screenshots showing the binding of CTCF, cohesin subunit (that is, RAD21) and ER α at *KCNK5* and surrounding gene promoters. **g, h**, qRT-PCR showing the mRNA and eRNA expression of *PGR* (**g**) and the expression of neighbouring genes (*ARHGAP42* and *TRPC6*) ($n = 3$) (**h**), in wild-type ($n = 2$) versus *PGR* promoter KO (*PGRp-KO*) ($n = 2$) independent isogenic cell clones of MCF-7 cells. Each bar represents an independent cell clone ($n = 3$ data points of technical replicates; representative of two independent experiments). **i**, Top, Hi-C contact matrix and ChIP-seq tracks of CTCF and p300 showing the topology of the chromosomal neighbourhood of the *PGR* locus. Bottom, UCSC browser screenshots showing CTCF, cohesin and ER α binding at *PGR* and surrounding gene promoters. Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's *t*-test.




Extended Data Fig. 3 | Cohesin knockdown affects gene and eRNA transcription through coordinating looping, without changing histone methylation. **a**, qRT-PCR results showing efficient knockdown of *RAD21* by siRNA, which does not affect the mRNA level of *ESR1* (encoding ER α) ($n = 3$ data points of technical replicates; representative of two independent experiments). **b**, **c**, qRT-PCR results showing that siRAD21 decreases the 17- β -oestradiol-induced activation of coding genes (**b**), but upregulates the eRNAs next to these genes (**c**) ($n = 3$ data points of technical replicates; representative of two independent experiments). **d**, MA plots showing the differential expression of RefSeq genes (left) or eRNAs (right) in cells with depletion of *RAD21* versus cells transfected with control siRNA (that is, siCTL). Red dots represent significantly changed genes or eRNAs (fold change > 1.5; FDR < 0.05). The purple bars indicate twofold change. **e**, UCSC genome browser screen shoots of GRO-seq results in siRAD21- versus siCTL-transfected MCF-7 cells at *KCNK5* (left) and *P2RY2* (right) loci; p300 ChIP-seq serves to indicate active enhancers. Yellow highlights denote gene regions; pink highlights denote enhancer regions. **f**, Diagram showing an oppositely regulated transcription of many eRNAs and mRNAs caused by disruption of E-P looping (that is, siRAD21). **g**, The tag density plots showed no significant difference of histone marks for promoters (H3K4me3 ChIP-seq) or enhancers (H3K4me1 ChIP-seq). qPCR data represent mean $\pm$ s.d.; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's t -test.

Red dots represent significantly changed genes or eRNAs (fold change > 1.5; FDR < 0.05). The purple bars indicate twofold change. **e**, UCSC genome browser screen shoots of GRO-seq results in siRAD21- versus siCTL-transfected MCF-7 cells at *KCNK5* (left) and *P2RY2* (right) loci; p300 ChIP-seq serves to indicate active enhancers. Yellow highlights denote gene regions; pink highlights denote enhancer regions. **f**, Diagram showing an oppositely regulated transcription of many eRNAs and mRNAs caused by disruption of E-P looping (that is, siRAD21). **g**, The tag density plots showed no significant difference of histone marks for promoters (H3K4me3 ChIP-seq) or enhancers (H3K4me1 ChIP-seq). qPCR data represent mean $\pm$ s.d.; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's t -test.

Extended Data Fig. 4 | Deletion of the *TFF1* promoter alters the transcription of the cognate *TFF1* enhancer and neighbouring genes.


a, Hi-C contact matrix in GM12878 cells (by Juicebox), CTCF-mediated chromatin loops (by ChIA-PET) and ChIP-seq tracks of CTCF and p300 in MCF-7 cells showing the topology (multiple contact domains) of the chromosomal neighbourhood of the *TFF1* locus. Yellow triangles in the Hi-C map denote contact domains; domains are numbered for simplicity. The contact domains containing genes analysed in **b** are highlighted in yellow. **b**, Box plots of gene expression outside of the hosting contact domain of *TFF1* generated by qRT-PCR in wild-type ($n = 6$) and *TFF1p-KO* ($n = 14$) isogenic clones of MCF-7 cells. The box plot centre lines represent medians; box limits indicate the 25th and 75th percentiles as determined by R software; and whiskers extend $1.5 \times \text{IQR}$ from the 25th and 75th percentiles. **c**, qRT-PCR results showing the expression of *TFF3* and *RSPH1* mRNA in wild-type MCF-7 cells as compared to *TFF1p-KO* cells

with two different gRNA pairs (*TFF1p-KO_gRNA1/2* and *TFF1p-KO_gRNA3/4*) ($n = 3$ data points of technical replicates; representative of three independent experiments). **d**, ChIP-qPCR data indicating the binding of RNA Pol II and p300 at *TFF3p* in wild-type cells as compared to *TFF1p-KO* cells ($n = 3$ data points of technical replicates; representative of two independent experiments). **e**, qRT-PCR results showing the expression of *TFF1e* and *TFF3m* after knockdown of *RAD21* by siRNA in wild-type MCF-7 cells versus *TFF1p-KO* cells ($n = 3$ data points of technical replicates; representative of two independent experiments). **f**, qRT-PCR results showing the expression *TFF1e*, *TFF1m* and *TFF3m* in wild-type MCF-7 cells versus cells with deletion of the *TFF1* enhancer (*TFF1e-KO*), *TFF1* promoter (*TFF1p-KO*) or both (*TFF1e/p-DKO*) ($n = 3$ data points of technical replicates; representative of three independent experiments). The diagram shows the *TFF1e* deletion in *TFF1p-KO* cells. Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's *t*-test.




Extended Data Fig. 5 | Epigenomic features of the enhancers and promoters that are putative looping pairs, as revealed by their opposite regulation after *siRAD21* treatment. **a**, ChIP-seq profile plots were generated using published data in MCF-7 cells (Supplementary Table 6), showing the differential enrichment of chromatin-looping-related structural factors, including CTCF, cohesin subunit (that is, RAD21), cohesin loading factor (that is, NIPBL) and Mediator subunit (that is, MED1) on the promoters that are downregulated by *siRAD21* treatment, as well as on enhancers for which eRNAs are increased by

siRAD21. Also refer to Extended Data Fig. 3. **b**, ChIP-seq profile plots showing the differential enrichment of factors that are relevant to enhancer function in MCF-7 cells, including those of p300, ER α , FOXA1 and GATA3. **c**, Diagram showing the non-stoichiometric distribution of chromatin structural factors and other transcription-related factors on functional E-P pairs. These E-P pairs denote RAD21-regulated enhancers (super-activated by *siRAD21*) and RAD21-dependent promoters (downregulated by *siRAD21*).




Extended Data Fig. 6 | Promoters exhibit a higher frequency and affinity of CTCF binding than enhancers. a, Percentages of promoters (P), enhancers (E) or random genomic regions (R) that contain a CTCF motif (left) or undergo CTCF binding (right) in MCF-7 cells (based on ENCODE data). The cartoons at the bottom indicate that promoters were further analysed for their features in **c**, on the basis of their distance to an adjacent enhancer. **b**, Similar to **a** but using GM12878 data. **c**, Percentage of MCF-7 promoters that contain a CTCF motif (left) or undergo CTCF binding (right) for three categories of promoters: promoters that have a super-enhancer, a typical enhancer or no enhancer in their 50-kb genomic proximity (cartoon at bottom of **a**). **d**, Similar to **c** but using GM12878 data. All *P* values in **a–e** were generated by two-sided Fisher's exact test. **e**, Top, diagram showing the design of constructs. The binding between GCN4 and ScFV will bring CTCF to the sites at which dCas9/gRNA

binds. Bottom, western blots showing the expression of scFv-CTCF-V5 (wild type or Y226A/F228A mutant) and dCas9-10xGCN4 after lentiviral transduction (uncropped images in Supplementary Fig. 1; representative of two independent experiments). **f**, ChIP-qPCR using a V5 antibody indicating comparable binding of V5-tagged wild-type CTCF or CTCF(Y226A/F228A) on *TFF3p* ($n = 3$ data points of technical replicates; representative of two independent experiments). Control gRNAs did not bring the V5-tagged CTCF to *TFF3p*. ChIP was performed with MCF-7 cells expressing dCas9-10xGCN4 and scFv-CTCF (wild type or Y226A/F228A mutant) as indicated. qPCR data in **f** represent mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, two-tailed Student's *t*-test. The *P* values were based on comparisons between the same-coloured bar (either wild-type or mutant CTCF) in TFF3p-gRNA1/gRNA2 versus CTL-gRNA1/gRNA2 conditions.



Extended Data Fig. 7 | Enhancer scanning and alternative models to interpret ERR and enhancer–promoter functional engagement. **a**, Model describing the process through which an activated enhancer scans its chromosomal neighbourhood inside a contact domain to functionally engage with its cognate promoter target. This enhancer scanning process involves promoter-bound CTCF (green ovals) and is compatible with cohesin-mediated extrusion (dashed lines) in wild-type cells, but it continues to operate in the absence of the cognate promoter (that is, P1) owing to deletion or disease mutation. A hypothetical ‘enhancer engagement’ time is depicted that reflects the relative amount of time in which the active enhancer engages with the neighbouring promoters, correlating with their expression levels. **b**, There are alternative, non-exclusive models that could be largely consistent with our results. For example, one of the other possibilities is a ‘positional effect’ model. In this model, an active enhancer may engage with its preferred promoter, and the two are retained in a transcription-associated E-P loop, quite probably in a nuclear environment such as an interchromatin granule⁵⁰, transcription factory or other perhaps phase-separated structure (pink coloured area). Upon functional loss (deletion, mutation or CTCF loss) of the original promoter, this ‘positional’ effect will follow the enhancer but it will engage with the secondary choices (P3 in the model; for example, *TFF3* as compared to *TFF1*, or *PAX5* as compared to *ZCCHC7*, or *TERT* as compared to *CLPTMIL*). But at a lower frequency or in a small percentage of single cells among a population, the enhancer does engage with other target promoters (such as P2, dotted black line). When the secondary choice is also lost, the entire enhancer will now select the third choice—for example, P2 in the model—to activate the promoter as its newest hierarchical choice. Overall, a few specific results from our study are in better support of the enhancer scanning model. First, it is consistent with the observation that for all single-cell colonies investigated (that is, 14 independent cell clones of *TFF1p*-KO in Extended Data Fig. 1), they consistently exhibited the highest expression of *TFF3* as the new hierarchical promoter choice, precluding models suggesting stochastic promoter choice. A positional effect model may predict that the enhancer and the target promoter are engaged so that the inhibition of one will reduce the other. This is true in

that deletion of *TFF1e* inhibited *TFF1p* (Extended Data Fig. 4f). However, CRISPRi of *TFF1p* resulted in higher activity of *TFF1e* and higher expression of *TFF3* (Fig. 3b). This data better support an enhancer scanning model—a dynamic enhancer–promoter interaction process inside the contact domain, in which *TFF1p* inhibition makes the ‘*TFF1e*-in-action’ preferentially interact with the next target in the hierarchy. Second, *TFF1e* exhibited quite broad chromatin interaction throughout the contact domain (for example, Fig. 1e, by 4C-seq), which is in accord with its ‘scanning’ behaviour. Furthermore, high-resolution Micro-C^{2,51} data showed that many ‘stripe loops’ formed in between enhancer and promoter at a finer scale (<50 kb or sometimes <10 kb), consistent with the suggestion that at least a subset of enhancers and promoters are actively scanning or extruding. The positional effect model is fundamentally compatible with the enhancer scanning model. The enhancer scanning model reflects a dynamic process in which the enhancer initiates target searching and it finds a major target as well as many additional minor targets. Were we to snapshot this dynamic process, then at every time point that an enhancer engages with one of its potential targets (for example, *TFF3p* or *TFF1p*), the target promoter would be repositioned closer to the enhancer. Our data suggest that the scanning process of the enhancer requires the cohesin complex, as its depletion by RNA interference reproduced many of the phenomena that are seen in promoter-knockout cells (for example, Fig. 1b, Extended Data Figs. 3, 4e). This is consistent with a proposed loop-extrusion model, rather analogous to the mechanism by which cohesin facilitates the formation of the larger TAD structures^{23,24,52}, requiring opposing CTCF motif orientation and high-affinity CTCF binding. However, it cannot be excluded that other chromatin remodellers, or Pol II itself, are the critical driver of enhancer scanning and E-P engagement. In support of this, inhibition of Pol II elongation partially reduced the promoter-centred stripe loops in mouse embryonic stem cells in a Micro-C study². Finally, it is noteworthy that the scanning concept that enhancers (either the DNA–protein complex as an entirety or specific transcription apparatus such as Pol II) travel along chromatin to reach target genes has been extensively discussed as one of the classic models to interpret enhancer activity^{22,53}.




Extended Data Fig. 8 | See next page for caption.

Extended Data Fig. 8 | CTCF binding on promoters, but not on enhancers. dictates enhancer–promoter choice. **a**, Bar graph showing the numbers of RAD21-regulated enhancers and promoters (same set used in Fig. 1b, Extended Data Fig. 5) that engage in chromatin loops in MCF-7 cells as discovered by analysing MCF-7 ChIA-PET data from ENCODE, as compared to their low incidence of looping in K562 ChIA-PET data. **b**, Coloured map showing that, among the functionally looped E-P pairs discovered in **a**, only certain percentages are bound by ER α , FOXA1 or CTCF by ChIP–seq. **c**, The numbers of looped E-P pairs that display various motif directionality of CTCF and FOXA1. The four categories below the plot describe the existence of CTCF or FOXA1 motifs on enhancers (E) or promoters (P); the colours indicate the combinations of motif directionality. For example, ‘CTCF (E) + CTCF (P)’ denotes the coincidence of CTCF motifs on both promoters and enhancers. There was no obvious orientation preference of binding motifs for CTCF. FOXA1 binds to functionally looped enhancers more frequently than CTCF

(Extended Data Figs. 5b, 8b), but it also exhibits no preferred motif orientation. **d**, Diagrams showing the position and directionality of CTCF-binding peaks and motifs in the four E-P pairs that we have extensively studied in this work. The colour motif sequence indicates directionality (black for left-pointing motifs, red for right-pointing motifs); the letters in each motif sequence are coloured to match the core CTCF motif nucleotides shown in the canonical CTCF motif by Jaspar (upper left logo); the motif scores after each motif sequence were calculated by the FIMO motif toolset. **e**, qRT–PCR results showing the expression of *TFF1* eRNA and mRNA in wild-type MCF-7 cells versus cells with deletion of the *TFF1* enhancer (*TFF1e-KO*), and MCF-7 cells with deletion of a CTCF peak inside the *TFF1* enhancer (*TFF1e/CTCF-KO*; Supplementary Table 1, Supplementary Fig. 3) ($n = 3$ data points of technical replicates; representative of two independent experiments). Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student’s *t*-test.


Extended Data Fig. 9 | Cancer mutations in gene promoter regions, and their link to oncogene activation through ERR. **a**, Plots showing the distribution of ICGC cancer whole-genome sequencing (WGS) mutations in the ± 5 kb regions near gene TSSs. The two numbers in each plot indicate the mutations in the ± 5 kb of TSSs, as well as the total numbers of mutations in that cancer type, respectively. Cancer type abbreviations can be found in the ICGC web portal (<http://icgc.org/>), and in the Methods. **b**, The percentage of genes containing ICGC annotated promoter mutations or deletions among all RefSeq genes (all gene promoters; total 27,502), oncogene promoters (OPs, from cosmic database; total 315) and neighbouring gene promoters within ± 200 kb of cosmic oncogenes (oncogene neighbouring promoters (ONPs); total 1,693). Hypergeometric tests were performed to compare the statistical significance of the enrichment of bearing a mutation or deletion for OPs or ONPs as compared to all RefSeq gene promoters (upon different frequency or size of mutations as indicated in the x axis). For each test, for example, comparing OPs to all RefGene promoters, $p[X > k]$ was calculated, which denotes a one-sided probability that a higher percentage of oncogene promoters bears mutations or deletions than the percentage of RefSeq promoters. The random variable X follows a hypergeometric distribution with parameters N (number of total RefSeq genes), K (number of total RefSeq genes bearing promoter mutations or deletions), n (number of total oncogenes or oncogene neighbours), k (number of total oncogenes or oncogene neighbours that bear promoter mutations or deletions). The five significant P values shown in the plot (from left to right) are 9.2×10^{-4} , 3.3×10^{-3} , 5.2×10^{-5} , 5.3×10^{-3} and 5.6×10^{-4} , respectively. $\text{mut} \geq 1$: any mutations being identified in ICGC; $\text{mut} > 1$: recurrent promoter mutations identified in more than 1 donors in ICGC release 28; $\text{del} < 100$ kb: genetic deletions with length smaller than 100 kb. **c**, Occurrence of somatic mutations and deletions at a selected list of ONPs that were included in our CRISPRi screening (see Fig. 3c). In this plot, the cancer cohorts (x axis) were ranked by the names of original cancer sites, and the y axis shows the ONP–OP gene pairs. Gene names before ‘->’ are those of ONPs, whereas those after ‘->’ are oncogenes. For example, ‘mutation: CLPTMIL -> TERT’ (the fourth row) indicates that *TERT* is an oncogene listed in COSMIC, and the gene promoter of

CLPTMIL was identified as an ONP, which contains somatic mutations in several cancer types. The dot size was scaled by the percentage of affected donors in each of that cancer type or cohort (SamplePerc), and dots of red colour show those samples with mutations, whereas blue dots show those samples with deletion (<100 kb) covering an ONP. The cancer type abbreviations can be found in the ICGC web portal (<http://icgc.org/>) and in the Methods. **d**, qRT-PCR results showing the expression of gene pairs, *PAX5-ZCCHC7*, *MYC-PVT1* and *SSI1L1-MTG2*. Each pair consisted of an oncogene (*PAX5*, *MYC* and *SSI1L1*) and its adjacent ONP genes (*ZCCHC7*, *PVT1* and *MTG2*). The ONP promoters were inhibited with specific sgRNAs in MCF-7 cells together with dCas9-KRAB (see the diagrams at the top) ($n = 3$ data points of technical replicates; representative of two independent experiments). **e, f**, Similar to Fig. 3d, these are results for two other ONP–OP loci after ONP deletion. The Hi-C contact matrix at the top shows the relative location of *PAX5-ZCCHC7* and *MYC-PVT1* gene pairs in a shared contact domain. qRT-PCR results below show the expression of ONP and OP gene mRNAs (two independent clones each) ($n = 3$ biological replicates). **g–i**, Oncoplots showing the landscapes of ICGC mutations located in three pairs of OPs as well as their ONPs at the *TERT-CLPTMIL* locus (**g**), the *PAX5-ZCCHC7* locus (**h**) and the *MYC-PVT1* locus (**i**). The percentages of mutations between 0.5–1% are denoted as 1%, and these between 0–0.5% are denoted as 0%. Each column in the oncoplot represents a cancer sample. The bar graphs with numbers (for example, 187, 101, 98) in the right side indicate the numbers of donor samples that contain mutations for that specific gene promoter. The labels on top of each panel indicate the total numbers of donors that contain mutations in any of the promoters in that specific locus, for example, 386 (6.27%) is the sum number of donors that contain mutations for any of the promoters shown in **g**. It is noteworthy that it remains a challenge to directly compare oncogene expression changes in tumour samples that carry rare noncoding mutations, because the rarity of matched RNA-seq and genotype (WGS) precludes a robust statistical analysis. qPCR data in **d–f** represent mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's t -test.



Extended Data Fig. 10 | Chromosomal interaction in three pairs of oncogene and oncogene-neighbourhood genes exhibiting an ERR-like phenotype. **a–c**, UCSC and WashU genome browser screenshots of ChIA-PET and GRO-seq results at the *PAX5–ZCCHC7* locus (**a**), *MYC–PVT1* locus (**b**) and *TERT–CLPTM1L* locus (**c**). The putative enhancers and promoters are highlighted in pink and in yellow, respectively. These cells show strong interactions between the eRNA (enhancer) and the original promoters (that is, *ZCCHC7p* in **a**, *PVT1p* in **b** and *CLPTM1Lp* in **c**). **d**, 4C-seq contact matrix and heat map plots showing the chromatin contacts with the *PAX5* promoter as

viewpoint, and its change in 293T cells with *ZCCHC7* promoter knockout. The viewpoint *PAX5* promoter is highlighted in yellow, and the neighbouring enhancer in pink. **e**, 4C-seq contact matrix and heat map plots showing the chromatin contacts with the *TERT* promoter as viewpoint, and its change in 293T cells with *CLPTM1L* promoter knockout. The viewpoint *TERT* promoter is highlighted in yellow, and the neighbouring enhancer in pink. **f**, The numbers of ICGC cancer mutations that reside in the motifs of CTCF, ERα, FOXA1 and GATA3, and random genomic regions near gene TSSs.



Extended Data Fig. 11 | GTEx data analysis identifies potential ERR gene pairs acting in human populations. **a**, Schematic overview of GTEx data-processing workflow to identify potential ERR events. Also see Supplementary Figs. 8–10 and associated notes. **b**, Histogram showing the number of potential ERR gene pairs that appear in one tissue or more than one. **c**, Histogram showing the number of potential ERR events distributed in each tissue. **d**, qRT-PCR results of CRISPRi showing the expression of *NUCKS1* and *RAB7L1* mRNAs with sgRNA-control (iCTL) or sgRNA specific for the putative enhancer (iEnhancer) in two clones of NPCs expressing dCas9-KRAB (see

enhancer location in Fig. 4c, d) ($n = 3$ data points of technical replicates; representative of two independent experiments). **e**, Published CTCF ChIP-seq screen shots in human-iPS-cell-derived neurons or neuronal cell lines (SK-N-SH) in the *NUCKS1* and *RAB7L1* promoters. The ChIP-seq data tracks were generated by the Cistrome data browser (<http://cistrome.org/>). The location of the three SNPs is shown. **f**, Sanger sequencing results of the *NUCKS1* promoter DNA, showing the allele bias of CTCF binding (comparison between the input DNA and ChIP DNA). Data are mean $\pm$ s.d.; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$, two-tailed Student's t -test.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a Confirmed

- ☐ ☒ The exact sample size (*n*) for each experimental group/condition, given as a discrete number and unit of measurement
- ☐ ☒ An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☐ ☒ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☒ ☐ A description of all covariates tested
- ☐ ☒ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☐ ☒ For null hypothesis testing, the test statistic (e.g. *F*, *t*, *r*) with confidence intervals, effect sizes, degrees of freedom and *P* value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's *d*, Pearson's *r*), indicating how they were calculated
- ☐ ☒ Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection	StepOnePlus™ Software v2.3 PXI 6 GeneSys software
Data analysis	Home-made codes are deposited in GitHub (https://github.com/wblilab-uth/ERR-project and https://github.com/zdz-lab/ERR). GraphPad Prism version 9 Juicebox 1.11.08 samtools 1.9 Bowtie 2 version 2.2.6 Bedtools version 2.25.0 HOMER version 4.8.2 MACS2 version 2.1.2 Illumina Basecalling Pipeline CASAVA 1.8.4 R/Bioconductor package version 3.6.1 The usages of software were described in the method section. All codes are available upon request. Data and Code availability information has been provided in the manuscript as a separate section.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

Code and Data availability statement was included in Methods Section.

Data and Code Availability: The datasets generated during and/or analyzed during the current study are available in the GEO (GSE115604) and SRA (PRJNA412021). For analyses of GTEx data and cancer mutations, codes are deposited in GitHub (<https://github.com/wblilab-uth/ERR-project> and <https://github.com/zdz-lab/ERR>). Other analyses of ChIP-Seq peaks, 4C-Seq or gene expression in the current study used standard bioinformatics tools and codes, which can be available upon request.

The association with traits and diseases were identified using genome wide association studies in the NHGRI-EBI GWAS Catalog(<https://www.ebi.ac.uk/gwas/>) and GWASdb v2(<https://doi.org/10.1093/nar/gkv1317>).

The GTEx data used for the analyses described in this manuscript were obtained from dbGap accession: phs000424.v7.p2.c1 (https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000424.v7.p2).

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☒ Life sciences ☐ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/authors/policies/ReportingSummary-flat.pdf](https://www.nature.com/authors/policies/ReportingSummary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	Sample size calculation was not performed. We determined the number of samples in each experiment as commonly accepted standards in the field. For qPCR and ChIP-qPCR, two or three biological replicates, each of three technical replicates was conducted. For key NGS data such as GRO-seq and 4C-seq, two to three biological replicates were conducted. We described these in the Figure Legends, Methods Section and Extended Data table 5.
Data exclusions	No data were excluded.
Replication	The data in this paper is highly replicable, as two laboratories conducted many experiments separately and reproduced all conclusions. We generated many promoter knockout isogenic cell clones to exclude single-cell clone-specific phenomenon. We performed two biological replicates for Extended Data Fig. 1e; 2a, b, d, e, g, h; 3a-c; 4d, e; 6f; 8e; 9d; 11d three biological replicates for Fig. 2c, d, f, g; 3b, d, f; 4e; Extended Data Fig. 1a, d; 4c, f; 9e, f; four biological replicates for Fig. 3g; 4f GRO-seq of RAD21 KD: 3 replicates, GRO-seq of TFF1pKO cells; 2 replicates, SMC1a, H3K4me3, H3K4me1 ChIP: n=1 but confirmed in +/- E2 signals. 4C-seq for TFF1e, TFF3p, KCNK5e, P2RY2e, NUCKS1p, RAB7L1p: 2 replicates
Randomization	This study does not involve work that requires random allocation. Most of the experiments are comparisons of isogenic control vs mutant cell line or control cells vs cells under specific treatment.
Blinding	The investigators were not blinded to data analysis (e.g. gene expression qPCR, GRO-seq or ChIP-seq). But for this current study, blinding is not relevant.

Reporting for specific materials, systems and methods

Materials & experimental systems

n/a	Involved in the study
☒	☐ Unique biological materials
☐	☒ Antibodies
☐	☒ Eukaryotic cell lines
☒	☐ Palaeontology
☒	☐ Animals and other organisms
☒	☐ Human research participants

Methods

n/a	Involved in the study
☐	☒ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging

Antibodies

Antibodies used

antibody information was provided in Method Section
 ERα Santa Cruz Sc543 ChIP and ChIP-seq
 p300 Santa Cruz Sc585 ChIP and ChIP-seq
 RNA Pol II Santa Cruz Sc-899 ChIP and ChIP-seq
 H3K4me3 Abcam Ab8580 ChIP-seq
 H3K4me1 Abcam Ab8895 ChIP-seq
 SMC1a Bethyl A300-055A ChIP-seq
 BrU agarose beads Santa Cruz Sc-32323 ac GRO-seq
 V5 Invitrogen R960-25 WB
 Cas9 Proteintech 26758-I-AP WB
 CTCF Millipore 07-729 ChIP and WB

Validation

All the antibodies used in this paper have been widely used and cited. The ChIP-seq pattern (e.g. H3K4me3 that mostly locates to promoters) also indicated that the the antibody is specific.
 ERα Santa Cruz <https://www.scbt.com/p/eralpha-antibody-hc-20>
 p300 Santa Cruz <https://www.scbt.com/p/p300-antibody-c-20>
 RNA Pol II Santa Cruz <http://datasheets.scbt.com/sc-899.pdf>
 H3K4me3 Abcam <https://www.abcam.com/histone-h3-tri-methyl-k4-antibody-chip-grade-ab8580.html>
 H3K4me1 Abcam <https://www.abcam.com/histone-h3-mono-methyl-k4-antibody-chip-grade-ab8895.html>
 SMC1a Bethyl <https://www.bethyl.com/product/A300-055A/SMC1+Antibody>
 BrU agarose beads Santa Cruz <https://www.scbt.com/p/brdu-antibody-iib5>
 V5 Invitrogen <https://www.thermofisher.com/antibody/product/V5-Tag-Antibody-Monoclonal/R960-25>
 Cas9 Proteintech <https://www.ptglab.com/products/Cas9-Antibody-26758-1-AP.htm>
 CTCF Millipore https://www.emdmillipore.com/US/en/product/Anti-CTCF-Antibody,MM_NF-07-729

Eukaryotic cell lines

Policy information about [cell lines](#)

Cell line source(s)

Cell lines, MCF-7, HEK293T, were from ATCC originally.
 Two hiPSC (iPSCORE_2_11 and iPSCORE_19_1) were obtained from iPSCORE at UCSD and were differentiated to NPCs.
 These are also described in the Method section.

Authentication

We purchase new batch of cells every 1-2 years from ATCC to ensure the authentication.

Mycoplasma contamination

Described in Methods. MCF-7 and HEK293T were examined for Mycoplasma contamination every 6-12 months.

Commonly misidentified lines
(See [ICLAC](#) register)

No commonly misidentified cell lines were used in our study.

Replicates

For key experiments, two to three biological replicates were conducted (e.g. GRO-seq for siRAD21 or promoter KO cells.)

Sequencing depth

ChIP-seq-SMC1a_EtOH 28,677,353
 ChIP-seq-SMC1a_E2 28,940,838
 ChIP-seq-H3K4me3_siCTL + E2 11,576,209
 ChIP-seq-H3K4me3_siRAD21 + E2 15,215,510
 ChIP-seq-H3K4me1_siCTL + E2 16,680,182
 ChIP-seq-H3K4me1_siRAD21 + E2 23,303,290
 ChIP-seq-H3K4me3_WT MCF7 + E2 17,872,206
 ChIP-seq-H3K4me3_TFF1p-KO MCF7 + E2 23,200,670
 GRO-seq-nascent RNA_siCTL + EtOH (repeat1) 6,863,150
 GRO-seq-nascent RNA_siCTL + E2 (repeat1) 7,667,228
 GRO-seq-nascent RNA_siRAD21 + EtOH (repeat1) 9,471,922
 GRO-seq-nascent RNA_siRAD21 + E2 (repeat1) 6,926,569
 GRO-seq-nascent RNA_siCTL + EtOH (repeat2) 12,624,935
 GRO-seq-nascent RNA_siCTL + E2 (repeat2) 13,276,916
 GRO-seq-nascent RNA_siRAD21 + EtOH (repeat2) 15,855,814
 GRO-seq-nascent RNA_siRAD21 + E2 (repeat2) 16,884,136
 GRO-seq-nascent RNA_siCTL + EtOH (repeat3) 8,771,184
 GRO-seq-nascent RNA_siCTL + E2 (repeat3) 7,815,340
 GRO-seq-nascent RNA_siRAD21 + EtOH(repeat3) 5,762,301
 GRO-seq-nascent RNA_siRAD21 + E2 (repeat3) 15,631,681
 GRO-seq-nascent RNA_Wildtype cell EtOH 9,675,923
 GRO-seq-nascent RNA_Wildtype cell E2 11,844,838
 GRO-seq-nascent RNA_TFF1p-KO (clone28) EtOH 11,633,638
 GRO-seq-nascent RNA_TFF1p-KO (clone28) E2 13,282,685
 GRO-seq-nascent RNA_TFF1p-KO (clone25) EtOH 20,015,942
 GRO-seq-nascent RNA_TFF1p-KO (clone25) E2 17,259,248
 4C-seq_TFF1e_viewpoint_wt_R1 4,155,458
 4C-seq_TFF1e_viewpoint_TFF1pKO_R1 3,460,929
 4C-seq_TFF1e_viewpoint_wt_R2 1,237,120
 4C-seq_TFF1e_viewpoint_TFF1pKO_R2 648,531
 4C-seq_TFF1e_viewpoint_wt_R3 6,651,744
 4C-seq_TFF1e_viewpoint_TFF1pKO_R3 7,251,685
 4C-seq_TFF3p_viewpoint_wt_R1 683,138
 4C-seq_TFF3p_viewpoint_TFF1pKO_R1 189,726
 4C-seq_TFF3p_viewpoint_wt_R2 2,996,729
 4C-seq_TFF3p_viewpoint_TFF1pKO_R2 2,800,472
 4C-seq_TFF3p_viewpoint_TFF1pKO_R3 5,424,601
 4C-seq_TFF1e_viewpoint_mcf7_sictrl_E21h_R1 4,111,503
 4C-seq_TFF1e_viewpoint_mcf7_siRAD21_E21h_R1 3,990,561
 4C-seq_TFF1e_viewpoint_mcf7_sictrl_E21h_R2 32,084,133
 4C-seq_TFF1e_viewpoint_mcf7_siRAD21_E21h_R2 26,919,259
 4C-seq_P2RY2e_viewpoint_mcf7_sictrl_E21h_R1 6,068,548
 4C-seq_P2RY2e_viewpoint_mcf7_siRAD21_E21h_R1 5,769,008
 4C-seq_P2RY2e_viewpoint_mcf7_sictrl_E21h_R2 583,828
 4C-seq_P2RY2e_viewpoint_mcf7_siRAD21_E21h_R2 754,506
 4C-seq_KCNK5e_viewpoint_mcf7_sictrl_E21h_R1 1,007,371
 4C-seq_KCNK5e_viewpoint_mcf7_siRAD21_E21h_R1 2,090,372
 4C-seq_KCNK5e_viewpoint_mcf7_sictrl_E21h_R2 185,863
 4C-seq_KCNK5e_viewpoint_mcf7_siRAD21_E21h_R2 82,854
 4C-seq_TERTp_viewpoint_293t_WT 2,756,635
 4C-seq_TERTp_viewpoint_293t_CLPTM1Lp-KO 2,156,282
 4C-seq_PAX5p_viewpoint_293t_WT 5,421,873
 4C-seq_PAX5p_viewpoint_293t_ZCCHC7p-KO 3,231,924
 4C-seq_nucks1p_viewpoint_alt_R1 34,713
 4C-seq_nucks1p_viewpoint_ref_R1 33,468
 4C-seq_nucks1p_viewpoint_alt_R2 37,309
 4C-seq_nucks1p_viewpoint_ref_R2 49,169
 4C-seq_RAB7L1p_viewpoint_alt_R1 18,388
 4C-seq_RAB7L1p_viewpoint_ref_R1 14,017
 4C-seq_RAB7L1p_viewpoint_alt_R2 253,592
 4C-seq_RAB7L1p_viewpoint_ref_R2 262,205

Antibodies

Described in Methods Section
 H3K4me3 Abcam Ab8580
 H3K4me1 Abcam Ab8895
 SMC1a Bethyl A300-055A
 BrU agarose beads Santa Cruz Sc-32323 ac

Peak calling parameters

MACS2 was used for peak calling using standard setting with q-value 0.01.

Data deposition

- ☒ Confirm that both raw and final processed data have been deposited in a public database such as [GEO](#).
- ☒ Confirm that you have deposited or provided access to graph files (e.g. BED files) for the called peaks.

Data access links

May remain private before publication.

GEO/GSE115604: <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE115604>
and SRA (PRJNA412021): <https://trace.ncbi.nlm.nih.gov/Traces/sra/?study=SRP118797>

Files in database submission

Provided in Extended Data Table 5

Genome browser session

(e.g. [UCSC](#))

https://genome.ucsc.edu/cgi-bin/hgTracks?hgS_doOtherUser=submit&hgS_otherUserName=wenbo%20li&hgS_otherUserSessionName=Oh_Paper_UCSC_2018

Data quality

ChIP-Seq reads were subjected to FASTQC for quality check.

Software

Samtools 1.9
Bowtie 2 version 2.2.6
Bedtools version 2.25.0
HOMER version 4.8.2
R/Bioconductor package version 3.6.1.